arXiv:cond-mat/0106076v1 [cond-mat.stat-mech] 5 Jun 2001

Statistical Theory for the Kardar–Parisi–Zhang Equation in $1+1$ dimension

A. A. Masoudi [a,e], F. Shahbazi [a,e], J. Davoudi [c,e]

and M. Reza Rahimi Tabar [b,d,e]

[a] *Dept. of Physics , Sharif University of Technology, P.O.Box 11365-9161, Tehran, Iran.*

[b] *CNRS UMR 6529, Observatoire de la Côte d'Azur, BP 4229, 06304 Nice Cedex 4, France,*

[c] *Max-Planck Institute for Complex Systems, Noethnitzer St.38, 01187 Dresden, Germany*

[d] *Dept. of Physics , Iran University of Science and Technology, Narmak, Tehran 16844, Iran.*

[e] *Institute for Studies in Theoretical Physics and Mathematics Tehran P.O.Box: 19395-5531, Iran,*

Abstract

The Kardar-Parisi-Zhang (KPZ) equation in 1+1 dimension dynamically develops sharply connected valley structures within which the height derivative *is not* continuous. There are two different regimes before and after creation of the sharp valleys. We develop a statistical theory for the KPZ equation in 1+1 dimension driven with a random forcing which is white in time and Gaussian correlated in space. A master equation is derived for the joint probability density function of height difference and height gradient $P(h-\bar{h}, \partial_x h, t)$ when the forcing correlation length is much smaller than the system size and much bigger than the typical sharp valley width. In the time scales before the creation of the sharp valleys we find the exact generating function of $h-\bar{h}$ and $\partial_x h$. Then we express the time scale when the sharp valleys develop, in terms of the forcing characteristics. In the stationary state, when the sharp valleys are fully developed, finite size corrections to the scaling laws of the structure functions $\langle (h-\bar{h})^n (\partial_x h)^m \rangle$ are also obtained. PACS: 05.70.Ln,68.35.Fx.

I. INTRODUCTION

There has been a great deal of recent work on the formation, growth and geometry of interfaces [1-5]. The dynamics of interfaces has turned out to be one of the most fascinating and at the same time challenging topics in theoretical non-equilibrium physics. There has been two principal approaches for theoretical analysis of such problems. The first is based on computer simulations of discrete models and often provides useful links between analytic theory and experiments. The second approach aims to describe the dynamical process by a stochastic differential equation. This procedure neglects the short length-scale details but provides a coarse-grained description of the interface (that is suitable for characterising the asymptotic scaling behaviour). Theoretical modelling of growth processes started with the work by Edwards and Wilkinson (EW) [6]. They suggested that one might describe the dynamics of the height fluctuations by a simple linear stochastic equation. Kardar–Parisi–Zhang (KPZ) [7], realized that there is a relevant term proportional to the square of the height gradient which represents a correction for lateral growth. Indeed the KPZ equation is a prototype model for a system in which the interface growth is subjected to a random external flux of particles. The randomness is described by an annealed random noise which mimics the random adsorption of molecules onto a surface. In the KPZ model (e.g in the 1+1 dimension), the surface height $h(x,t)$ on the top of location x of 1-dimensional substrate satisfies a stochastic random equation,

$$\frac{\partial h}{\partial t} - \frac{\alpha}{2}(\partial_x h)^2 = \nu \partial_x^2 h + f(x,t), \tag{1}$$

where $\alpha \geq 0$ and f is a zero-mean, statistically homogeneous, white in time and Gaussian process with covariance

$$\langle f(x,t) f(x',t') \rangle = 2D_0 \delta(t-t') D(x-x'), \tag{2}$$

where,

$$D(x-x') = \frac{1}{\sqrt{\pi}\sigma} \exp(-\frac{(x-x')^2}{\sigma^2}), \tag{3}$$

and σ is the variance of $D(x-x')$. Typically the correlation of forcing is considered as delta function for mimicking the short range correlation. We regularise the delta function correlation by a Gaussian function. When the variance σ is much less than the system size we would expect that the model would represent a short-range correlated forcing. So we would stress that our calculations are done for finite $\sigma \ll L$, where L is the system size. The average force on the interface is unimportant and may be removed from the equation of motion. Every term in the eq.(1) involves a specific physical phenomenon contributing to the surface evolution. The parameters ν, α and D_0 (and σ) are describing the surface diffusive relaxation, non-linear lateral growth and the effective noise strength, respectively.

We consider a substrate of size L and define the mean height of growing film and its roughness w by,

$$\bar{h}(L,t) = \frac{1}{L}\int_{-L/2}^{L/2} dx h(x,t), \tag{4}$$

$$w(L,t) = (\langle (h-\bar{h})^2 \rangle)^{1/2}, \tag{5}$$

where $\langle \cdots \rangle$ denotes an averaging over different realizations of the noise (samples). Starting from a flat interface (one of the possible initial conditions), it was conjectured by Family and Vicsek [8] that a scaling of space by factor b and of time by a factor b^z (z is the dynamical scaling exponent), re-scales the roughness w by factor b^χ as follows,

$$w(bL, b^z t) = b^\chi w(L,t), \tag{6}$$

which implies that

$$w(L,t) = L^\chi f(\frac{t}{L^z}). \tag{7}$$

If for large t and fixed L ($\frac{t}{L^z} \to \infty$), w saturates then $f(x) \to const.$ as $x \to \infty$. However, for fixed large L and $1 << t << L^z$, one expects that correlations of the height fluctuation are set up only within a distance $t^{1/z}$ and thus must be independent of L. This implies that for $x << 1$, $f(x) \sim x^\beta$ with $\beta = \frac{\chi}{z}$. Thus dynamic scaling postulates that [9],

$$w(L,t) \sim t^{\beta} \qquad 1 << t << L^{z}$$

$$\sim L^{\chi} \qquad t >> L^{z} \tag{8}$$

The roughness exponent χ and the dynamic exponent z characterise the self-affine geometry of the surface and its dynamics, respectively. Several time-regimes can be distinguished in the time evolution of the surface roughness. They can be summarised as follows: for very early times , the noise term dominates since its contribution to the equation grows as the square root of time. In this time-regime, the surface roughness grows as $w(t) \sim t^{1/2}$. For intermediate times, the linear term has the main contribution. The linear case $(\alpha = 0)$ is the Edwards-Wilkinson model for which one can easily find that the surface roughness behaves as $w(t) \sim t^{\beta_0}$, where the value of β_0 depends on dimension of substrate($\beta_0 = \frac{2-d}{4}$ for d-dimensional surface). For later times, the contribution of the relevant non-linear term becomes a dominant one and the surface roughness growth is characterised by the behaviour $w(t) \sim t^{\beta}$. For very late times and finite substrate-length L, the roughness saturates to the value $w(t \to \infty, L) \sim L^{\chi}$. Of course, in an experiment or in a numerical simulation the transition between the different regimes is not sharp and different crossover behaviours can be observed. Galilean invariance implies the relation $\chi + z = 2$ independent of dimension [10-11]. It means that there is only one independent exponent in the KPZ dynamics. In the one-dimensional substrates a fluctuation-dissipation theorem yields exactly $z = \frac{3}{2}$, $\chi = \frac{1}{2}$, $\beta = \frac{1}{3}$ [12]. In contrast to 1-dimension, the case $d \geq 2$ can be only attacked by approximative field-theoretic perturbative expansions [13-16]. It is well-known that effective coupling constant for the KPZ equation is $g = \frac{2\alpha^2 D_0}{\nu^3}$. Phase diagram information extracted from the renormalisation group flow indicates that $d = 2$ plays the role of a lower critical dimension. For $d \leq 2$, the Gaussian fixed point $(\alpha = 0)$ is infrared-unstable, and there is a crossover to the stable strong coupling fixed point. For $d > 2$, a third fixed point exists, which represents the roughening transition. It is unstable and appears between the Gaussian and strong coupling fixed points which are now both stable. Only the critical indices of the strong-coupling regime ($g \to \infty$ or $\nu \to 0$) are known in 1+1 dimensions and their values in higher dimensions as well as properties of the roughening transition have been known only numerically [17-23], and the various approximation schemes [24-32].

The theoretical richness of the KPZ model is partly due to close relationships with other areas of statistical physics. It is shown that there is a mapping between the equilibrium statistical mechanics of a two dimensional smectic-A liquid crystal onto the non-equilibrium dynamics of the (1+1)- dimensional stochastic KPZ equation [33]. It has been shown in [34] that, one can map the kinetics of the annihilation process $A + B \to 0$ with driven diffusion onto the (1+1)-dimensional KPZ equation. Also the KPZ equation is closely related to the dynamics of a sine-Gordon chain [35], the driven-diffusion equation [36-37], high T_c-superconductor [38] and directed paths in the random media [39-52] and charge density waves [53], dislocations in disordered solids [3], formation of large-scale structure in the universe [54-57] , Burgers turbulence [58-85,90] and etc.

As mentioned the main difficulty with the KPZ equation is that it is controlled, in all dimensions, by a strong disorder (or strong coupling) fixed point and efficient tools are missing to calculate the exponents and other universal properties e.g. scaling functions, amplitudes, etc. Despite the fact that in one dimension, the exponents are known, but many properties, including the probability density function (PDF) of the height of a growing interface have been so far measured only in numerical simulations. Recently, it is shown that for one particular model of the KPZ class, the asymmetric exclusion process (ASEP), the whole distribution of displacement of particles could be calculated for a finite geometry by Bethe ansatz [86-87]. As discussed in [86-87], the PDF of displacement of particles is related to the PDF of height differences in the growth process. It is proved in [86-87] that the PDF of height differences has the following asymptotic:

$$\begin{aligned} P(\frac{h-\bar{h}}{w}) \quad &\sim \quad \exp(-Ay^{5/2}) \qquad y \to +\infty, \\ &\sim \quad \exp(-B|y|^{3/2}) \qquad y \to -\infty, \end{aligned} \tag{9}$$

where $y = \frac{h-\bar{h}}{w}$ and w is the variance of the height fluctuation about its average. This result is in agreement with the numerical experiments [88].

In this paper we are interested in the statistical properties of the KPZ equation in the strong coupling limit ($\nu \to 0$). The limit is singular,i.e. through which the surface develops sharp valleys. So starting with a flat surface after a finite time scale, t_c, the sharp valley singularities are dynamically developed. In the singular points (sharp valleys) spatial derivative of the $h(x,t)$ is not continuous. Hence the limit of $\nu \to 0$ is not singular for $t < t_c$, and we can ignore the diffusion term while after developing the singularities the diffusion term has finite contribution in the PDF of height fluctuations. Inspired by the methods proposed, recently in the works done by Weinan E and Vanden Eijnden [73], we develop a statistical method to describe the moments of height difference and height gradient of height field $h(x,t)$. We derive a master equation for joint probability density function (PDF) of the height difference $(h-\bar{h})$ and height gradient $\partial_x h$, $P((h-\bar{h}), \partial_x h)$ for given g or diffusion constant ν. We will consider two different time scales in the limit of $\nu \to 0$. (i) Early stages before developing the sharp valley singularities and (ii) established stationary state comprising with fully developed sharp valley singularities. In the regime (i) ignoring the relaxation term in the equation of the joint PDF when $\nu \to 0$, we determine the exact generating function of joint moments of height and height gradient fields. Realisability condition for the resulting joint PDF sheds light on the time scale of the sharp valley formation. In contrary the limit $\nu \to 0$ is singular in the regime (ii) leading to an unclosed term (relaxation term) in the PDF equation. However we show that the unclosed term can be expressed in terms of statistics of some quantities defined on the singularities (sharp valleys). Identifying each sharp valley in position y_0 with three quantities, namely the gradient of h in the position y_{0^+}, y_{0^-} and its height from the $\bar{h}$, we determine the dynamics of these quantities. In the both regimes all the moments, $\langle (h-\bar{h})^n (\partial_x h)^m \rangle$, for given n and m are found. In the regime (ii) we will prove that in leading order, when $L \to \infty$, fluctuation of the height field is not intermittent and succeed to give the analytic form of amplitudes of the all structure functions. Besides, the scaling behaviour and the amplitudes of all the correction terms due to the finite size effect are calculated.

The paper is organised as follows; in section two, we derive the master equation for the joint PDF of height differences and height gradients for given diffusion ν. We convert the height PDF,i.e. $P(h-\bar{h},t)$, evolution equation consequently to a Fokker-Planck equation for an arbitrary given diffusion constant. In section three we will consider the limit of $\nu \to 0$ of the master equation in the time scales that there are no any singularities in the surface (before developing the sharp valleys). We determine the exact and explicit expression of the generating function for the moments $\langle (h-\bar{h})^n (\partial_x h)^m \rangle$ for given n and m. In section 4 we consider the master equation in the limit $\nu \to 0$ and consequently when the singularity are fully developed. In this regime the relaxation term has finite contribution in the master equation. Using the methods introduced in [73], we prove that the unclosed term can be written in terms of quantities which are defined on sharp valleys where $\partial_x h$ is discontinuous. Also in this section we determine the relation between the density of sharp valleys and the forcing variance $k_{xx}(0)$ in detail. In section 5, we derive the moments of height fluctuation in the stationary state and show that the PDF of $(h-\bar{h})$ is strongly asymmetric and prove that in to leading order the n-th moments of $(h-\bar{h})$, i.e. $\langle (h-\bar{h})^n \rangle$, can be written in terms of the second order moment of height fluctuation in a non-intermittent way. We determine the amplitudes of the all of moments and show that the amplitudes of moments $\langle (h-\bar{h})^n (\partial_x h)^m \rangle$ can be written in terms of characteristics of singularities. We also derive the finite size effect on the moments of height differences and determine the amplitudes of all correction terms. In section 6, we derive the PDF of quantities which characterise the singularity, and therefore find the equation for their evolution.

II. THE MASTER EQUATION FOR HEIGHT DIFFERENCE AND HEIGHT GRADIENT

In this section we consider the $1+1-$dimensional KPZ equation and derive the master equation to describe the joint-PDF of height-difference and height-gradient i.e. $P(h-\bar{h}, \partial_x h)$, for given ν and α. It is shown that the equation for the joint PDF is not closed due to the *linear* term $\nu\partial^2 h$. The PDF of height difference is related to the joint PDF, $P(h-\bar{h}, \partial_x h, t)$ by the relation $P(h-\bar{h}, t) = \int_{-\infty}^{\infty} P(h-\bar{h}, \partial_x h, t) d(\partial_x h)$. We show that $P(h-\bar{h}, t)$ satisfies a Fokker-Planck equation and write down the explicit expression of drift and diffusion coefficient $D^{(1)}$ and $D^{(2)}$. It is shown that the drift and the diffusion coefficients can be written in terms of the conditional average of $\langle(\partial_x h)^2 | h-\bar{h}\rangle$.

We consider a one-dimensional line of length L and a surface of height $h(x,t)$, and gradient $\partial_x h(x,t)$ at time t. The 1+1 dimensional KPZ equation governed over height field $h(x,t)$ is defined in eq.(1) while $u(x,t) = -\partial_x h(x,t)$ is a solution of the the so called Burgers equation as,

$$u_t + \alpha u u_x = \nu u_{xx} - f_x(x,t), \tag{10}$$

where the covariance of f is given by eqs.(2) and (3). To investigate the statistical properties of eqs.(10) and (1), let us define the generating function $Z(\lambda, \mu, x, t)$ as:

$$Z(\lambda, \mu, x, t) = \langle \exp(-i\lambda(h(x,t)-\bar{h}) - i\mu u(x,t))\rangle. \tag{11}$$

It follows from eqs.(10) and (1) that generating function Z is a solution of the following equation,

$$Z_t = i\gamma(t)\lambda Z - i\lambda\frac{\alpha}{2}Z_{\mu\mu} - \lambda^2 k(0) Z + \frac{i\alpha\lambda + \nu\lambda^2}{\mu}Z_\mu - \frac{\nu\lambda}{\mu}Z_x$$

$$-i\alpha\mu(\frac{Z_x}{\mu})_\mu + \mu^2 k_{xx}(0) Z - i\mu\nu\langle u_{xx}\exp(-i\lambda(h-\bar{h}) - i\mu u)\rangle, \tag{12}$$

where $k(x-x') = 2D_0 D(x-x')$, $\gamma(t) = \bar{h}_t$, $k(0) = \frac{D_0}{\sqrt{\pi}\sigma}$ and $k_{xx}(0) = -\frac{2D_0}{\sqrt{\pi}\sigma^3}$. To derive eq.(12) we have used the following identities:

$$\langle h_{xx} \exp\left(-i\lambda(h(x,t)-\bar{h}) - i\mu u(x,t)\right)\rangle = \frac{-i}{\mu}\{Z_x + \lambda\partial_\mu Z\}, \tag{13}$$

$$\langle f(x,t) \exp\left(-i\lambda(h(x,t)-\bar{h}) - i\mu u(x,t)\right)\rangle = -i\lambda k(0) Z, \tag{14}$$

and

$$\langle f_x(x,t) \exp\left(-i\lambda(h(x,t)-\bar{h}) - i\mu u(x,t)\right)\rangle = -i\mu k_{xx}(0) Z, \tag{15}$$

where we have used the fact that $D_x(0) = 0$. This is evident that due to the last term in the eq.(12), it is not closed. Assuming statistical homogeneity ($Z_x = 0$) we have,

$$\begin{aligned} -i\mu Z_t &= \gamma\lambda\mu Z - \frac{\alpha}{2}\lambda\mu Z_{\mu\mu} + i\lambda^2\mu k(0) Z - i\mu^3 k_{xx}(0) Z \\ &\quad - i(\nu\lambda^2 + i\alpha\lambda) Z_\mu - \mu^2\nu\langle u_{xx}\exp(-i\lambda\tilde{h}(x,t) - i\mu u(x,t))\rangle, \end{aligned} \tag{16}$$

where $\tilde{h}(x,t) = h(x,t) - \bar{h}$. Defining $P(\tilde{h},u,t)$ as the joint probability density function of $\tilde{h}$ and u, one can construct $P(\tilde{h},u,t)$ in terms of generating function Z as,

$$P(\tilde{h},u,t) = \int\int \frac{d\lambda}{2\pi}\frac{d\mu}{2\pi} \exp(i\lambda\tilde{h} + i\mu u) Z(\lambda,\mu,t). \tag{17}$$

It follows from eqs.(17) and (16) that $P(\tilde{h},u,t)$ satisfies the following equation,

$$\begin{aligned} -P_{ut} &= -\gamma P_{\tilde{h}u} - \frac{\alpha}{2}(u^2 P)_{\tilde{h},u} - \alpha(uP)_{\tilde{h}} - k(0) P_{\tilde{h}\tilde{h}u} + k_{xx}(0) P_{uuu} - \nu(uP)_{\tilde{h}\tilde{h}} \\ &\quad - \nu\int\int \frac{d\lambda}{2\pi}\frac{d\mu}{2\pi}\exp(i\lambda\tilde{h} + i\mu u)\mu^2\langle u_{xx}\exp(-i\lambda\tilde{h}(x,t) - i\mu u(x,t))\rangle. \end{aligned} \tag{18}$$

Now we can rewrite the last term in eq.(18) as following,

$$\begin{aligned} &\nu\int\int \frac{d\lambda}{2\pi}\frac{d\mu}{2\pi}\exp(i\lambda\tilde{h} + i\mu u)\mu^2\langle u_{xx}(x,t)\exp(-i\lambda\tilde{h}(x,t) - i\mu u(x,t))\rangle \\ &= -\nu\langle u_{xx}(x,t)\delta(\tilde{h}(x,t) - \tilde{h})\delta(u(x,t) - u)\rangle \\ &= -\nu\{\langle u_{xx}|u,\tilde{h}\rangle P(u,\tilde{h},t)\}_{uu}. \end{aligned} \tag{19}$$

where $\langle\cdot|u,\tilde{h}\rangle$ denotes the average conditional on a given $u,\tilde{h}$. Therefore using the eq.(19) it follows that the $P(\tilde{h},u,t)$ satisfies the following,

$$-P_{ut} = -\gamma P_{\tilde{h}u} - \frac{\alpha}{2}(u^2P)_{\tilde{h},u} - \alpha(uP)_{\tilde{h}} - k(0)P_{\tilde{h}\tilde{h}u} + k_{xx}(0)P_{uuu} - \nu(uP)_{\tilde{h}\tilde{h}} +$$

$$+\nu\{\langle u_{xx}|u,\tilde{h}\rangle P(u,\tilde{h},t)\}_{uu}. \tag{20}$$

This equation is exact for a given ν and as it is clear the fingerprints of diffusion term is giving rise again to an unclosed equation for $P(\tilde{h},u,t)$. Accessing to the functional form of the conditional averaging $\langle u_{xx}|u,\tilde{h}\rangle$ is one of the major difficulties in the formulation. From eq.(18) we see that $P(u,\tilde{h}) = P(-u,\tilde{h})$ which results to,

$$\langle u|\tilde{h}\rangle = 0, \tag{21}$$

for any given ν. In fact for a given h the average of height gradient, u, is consequently zero. The identity proposed by eq.(21) is not restricted to any limiting asymptotic and is true in all regimes of the dynamical evolution of the surface. Also eq.(20) allows us to determine a dynamical equation for $P(h-\bar{h})$. Doing so, we multiply the eq.(20) to u an integrate over u from $-\infty$ to $+\infty$, from which we get,

$$\partial_t P(\tilde{h},t) = \frac{\partial}{\partial\tilde{h}}\{(\frac{\alpha}{2}(\langle u^2\rangle - \langle u^2|\tilde{h}\rangle))P(\tilde{h},t)\}$$

$$+\frac{\partial^2}{\partial\tilde{h}^2}\{(k(0) - \nu\langle u^2|\tilde{h}\rangle)P(\tilde{h},t)\}, \tag{22}$$

where $\tilde{h} = h - \bar{h}$ and the relation $\gamma = \frac{\alpha}{2}\langle u^2\rangle$ is used. This is a Fokker-Planck (FP) equation, describing the time evolution of $P(\tilde{h},t)$. The drift coefficient in the FP equation is,

$$D^{(1)} = -\frac{\alpha}{2}(\langle u^2\rangle - \langle u^2|\tilde{h}\rangle), \tag{23}$$

and the diffusion coefficient reads,

$$D^{(2)} = k(0) - \nu\langle u^2|\tilde{h}\rangle. \tag{24}$$

It is evident that aiming to obtain $P(h-\tilde{h})$, one should know the conditional average $\langle u^2|\tilde{h}\rangle$. The equation has the following stationary solution:

$$P_{stat.}(\tilde{h}) = \frac{N}{D^{(2)}} \exp\{\int_{\tilde{h}_0}^{\tilde{h}} d\tilde{h}' D^{(1)}(\tilde{h}')/D^{(2)}(\tilde{h}')\} \tag{25}$$

Where N is the normalisation coefficient. Therefore to derive the moments of hight difference $h-\bar{h}$ i.e. $\langle (h-\bar{h})^n\rangle$, we need the conditional averaging $\langle u^2|\tilde{h}\rangle$. The simplified picture given by this equation is indicating that all the knowledge one would know in order to obtain the behaviour of PDF is buried in the functional form of one conditional average , i.e. $\langle u^2|\tilde{h}\rangle$. Although simple but it is clear that the conditional average $\langle u^2|\tilde{h}\rangle$ would have a non-trivial dependence on ν and L in the limit of $\nu \to 0$. Instead of following this strategy however in the next section we follow another direct way of extracting the moments of height difference $(h-\bar{h})$ in the strong coupling limit , i.e. $\nu \to 0$.

III. THE JOINT CORRELATIONS OF HEIGHT DIFFERENCE AND HEIGHT GRADIENT BEFORE SHARP VALLEY FORMATION

When σ is finite, the very existence of the non-linear term in the KPZ equation leads to the development of the sharp valley singularities in a *finite time* and in the strong coupling limit($\nu \to 0$). In one dimension system is already in the strong coupling regime so starting from any finite value of ν in large time system develops sharp valley singularities, Fig(2). Therefore one would distinguish between different time regimes before and after the sharp valley formation. Starting from a flat initial condition, i.e. $h(x,0)=0, u(x,0)=0$, which its evolution is given by inviscid KPZ equation, we know that after a finite time the derivative of function $h(x,t)$ becomes singular. After this time scale the diffusion term is important, but we can neglect that before appearance of the singularities. So the equation governing the the evolution of generating function, $Z(\mu,\lambda,t)$ before the creation of the sharp valleys is given by,

$$Z_t = i\gamma(t)\lambda Z - i\lambda\frac{\alpha}{2}Z_{\mu\mu} - \lambda^2 k(0)Z + \frac{i\alpha\lambda}{\mu}Z_\mu - i\alpha\mu(\frac{Z_x}{\mu})_\mu + \mu^2 k_{xx}(0)Z, \tag{26}$$

where we have assumed the statistical homogeneity ($Z_x = 0$). Now we need the γ which is given by $\gamma = \bar{h}_t$. So invoking to the eq.(10) we get,

$$\gamma(t) = \bar{h}_t = \frac{\alpha}{2}\langle u^2 \rangle. \tag{27}$$

To evaluate $\langle u^2 \rangle$ we set $\lambda = 0$ in the eq.(26), and find,

$$Z_t = \mu^2 k_{xx}(0) Z, \tag{28}$$

which considering $Z(\mu, 0) = 1$ as initial condition, its solution is,

$$Z(\mu, t) = \exp(\mu^2 k_{xx}(0) t). \tag{29}$$

On the other hand by definition we have $\langle u^2 \rangle = -(\frac{\partial^2 Z(\mu,t)}{\partial \mu^2})$. So before creation of the singularities the second moment of height gradient behave as:

$$\langle u^2 \rangle = -2k_{xx}(0)t, \tag{30}$$

so consequently,

$$\gamma(t) = -\alpha k_{xx}(0)t. \tag{31}$$

Inserting eq.(31) in the equation (26) results to,

$$\frac{\partial}{\partial t} Z(\mu, \lambda, t) = -i\alpha\lambda \frac{\partial^2}{\partial \mu^2} Z(\mu, \lambda, t) + i\alpha \frac{\lambda}{\mu} \frac{\partial}{\partial \mu} Z(\mu, \lambda, t)$$

$$+(\mu^2 k_{xx}(0) - i\alpha k_{xx}(0) t \lambda - \lambda^2 k(0)) Z(\mu, \lambda, t). \tag{32}$$

We solve the eq.(32) with the initial condition $Z(\mu, \lambda, 0)$=1, from which by expanding the generating function in powers of λ and μ we can obtain the moments $\langle (h - \bar{h})^n \rangle$,$\langle u^n \rangle$ and $\langle (h - \bar{h})^n u^m \rangle$. We change the variable μ to $y = \mu^2$, so the eq.(32) converts to the following equation,

$$\frac{\partial}{\partial t} Z(y, \lambda, t) = -2i\alpha\lambda y \frac{\partial^2}{\partial y^2} Z(y, \lambda, t) + i\alpha\lambda \frac{\partial}{\partial y} Z(y, \lambda, t)$$

$$+(yk_{xx}(0) - i\alpha k_{xx}(0)t\lambda - \lambda^2 k(0))Z(y,\lambda,t). \tag{33}$$

Introducing the Fourier transform of $Z(\mu,\lambda,t)$ with respect to y as $Q(q,\lambda,t)$, it is simple to get the following evolution equation satisfied by the Fourier transform,

$$\begin{aligned}\frac{\partial}{\partial t}Q(q,\lambda,t) = 2\alpha\lambda q^2 \frac{\partial}{\partial q}Q(q,\lambda,t) + 5\alpha\lambda q Q(q,\lambda,t) - ik_{xx}(0)\frac{\partial}{\partial q}Q(q,\lambda,t) + \\ -i\alpha k_{xx}(0)t\lambda Q(q,\lambda,t) - \lambda^2 k(0)Q(q,\lambda,t),\end{aligned} \tag{34}$$

with the initial condition

$$Q(q,\lambda,0) = \frac{1}{2\pi}\int e^{iyq}dy = \delta(q). \tag{35}$$

The eq.(34) is a first order partial differential equation which can be solved by the method of characteristics. The general solution of eq. (34) is written as,

$$\begin{aligned}Q(q,\lambda,t) = g\left(\lambda, \frac{1}{2}\frac{2tk_{xx}(0)\alpha\lambda + \sqrt{-2ik_{xx}(0)\alpha\lambda}\tanh^{-1}(q\sqrt{-\frac{2i\alpha\lambda}{k_{xx}(0)}})}{k_{xx}(0)\alpha\lambda}\right) \times \\ \times \exp\Bigg\{ -1/2\int_0^q \Big(\frac{10\alpha\lambda s + i\tanh^{-1}(s\sqrt{\frac{-2i\alpha\lambda}{k_{xx}(0)}})\sqrt{-2ik_{xx}(0)\alpha\lambda}}{2\alpha\lambda s^2 - ik_{xx}(0)} \\ \frac{+2i\alpha k_{xx}(0)t\lambda - 2\lambda^2 k(0) - i\tanh^{-1}(q\sqrt{\frac{-2i\alpha\lambda}{k_{xx}(0)}})\sqrt{-2ik_{xx}(0)\alpha\lambda}}{2\alpha\lambda s^2 - ik_{xx}(0)}\Big)ds\Bigg\},\end{aligned} \tag{36}$$

where g is an arbitrary function of its arguments. Imposing the initial condition, given in eq.(35), and introducing ω as,

$$\omega = \frac{1}{2}\frac{\tanh^{-1}(q\sqrt{\frac{-2i\alpha\lambda}{k_{xx}(0}})\sqrt{-2ik_{xx}(0)\alpha\lambda}}{k_{xx}(0)\alpha\lambda}, \tag{37}$$

we obtain,

$$\begin{aligned}g(\lambda,\omega) = \delta(-\frac{1}{2}\sqrt{2}\sqrt{\frac{ik_{xx}(0)}{\alpha\lambda}}\tanh(\sqrt{2ik_{xx}(0)\alpha\lambda}\omega))\exp[1/2\int_0^{-\frac{1}{2}\sqrt{2}\sqrt{\frac{ik_{xx}(0)}{\alpha\lambda}}\tanh(\sqrt{2ik_{xx}(0)\alpha\lambda}\omega)} \\ (\frac{10\alpha\lambda s + i\tanh^{-1}[s\sqrt{\frac{-2i\alpha\lambda}{k_{xx}(0)}}]\sqrt{-2ik_{xx}(0)\alpha\lambda} - 2i\omega k_{xx}(0)\alpha\lambda - 2\lambda^2 k(0)}{2\alpha\lambda s^2 - ik_{xx}(0)})ds],\end{aligned} \tag{38}$$

from which $Q(q,\lambda,t)$ is obtained as,

$$Q(q,\lambda,t) = g(\lambda, t+\omega)\exp[-1/2\int_0^q \frac{10\alpha\lambda s + i\tanh^{-1}[s\sqrt{\frac{-2i\alpha\lambda}{k_{xx}(0)}}]\sqrt{-2ik_{xx}(0)\alpha\lambda}}{2\alpha\lambda s^2 - ik_{xx}(0)}$$
$$\frac{-2i\alpha\lambda k_{xx}(0)t - i\tanh^{-1}[q\sqrt{\frac{-2i\alpha\lambda}{k_{xx}(0)}}]\sqrt{-2ik_{xx}(0)\alpha\lambda} - 2\lambda^2 k(0)}{2\alpha\lambda s^2 - ik_{xx}(0)} ds]. \tag{39}$$

Inverse Fourier transforming of the solution in eq.(39) is straightforward, so after switching to variable μ we get the following solution for $Z(\mu,\lambda,t)$:

$$Z(\mu,\lambda,t) = \left(1-\tanh^2(\sqrt{2ik_{xx}(0)\alpha\lambda}t)\right)\exp\Big\{-\frac{5}{8}\ln(1-\tanh^4(\sqrt{2ik_{xx}(0)\alpha\lambda}t))$$
$$+\frac{5}{4}\tanh^{-1}(\tanh^2(\sqrt{2ik_{xx}(0)\alpha\lambda}t)) - \lambda^2 k(0)t$$
$$-\frac{1}{16}\ln^2(\frac{1-\tanh(\sqrt{2ik_{xx}(0)\alpha\lambda}t)}{1+\tanh(\sqrt{2ik_{xx}(0)\alpha\lambda}t)}) - \frac{1}{2}i\mu^2\sqrt{\frac{2ik_{xx}(0)}{\alpha\lambda}}\tanh(\sqrt{2ik_{xx}(0)\alpha\lambda}t)\Big\}. \tag{40}$$

Since we are interested in moments $\langle(h-\bar{h})^n\rangle$, so setting $\mu = 0$ in (40) and expanding the generating function in power of λ we can obtain them all. For example expanding up to $O(\lambda^6)$ it is easy to see that the first sixth order of moments are behaving as following,

$$\langle(h-\bar{h})^2\rangle = -\frac{1}{3}t(k_{xx}(0)^2\alpha^2 t^3 - 6k(0)), \tag{41}$$
$$\langle(h-\bar{h})^3\rangle = -\frac{24}{45}k_{xx}(0)^3\alpha^3 t^6, \tag{42}$$
$$\langle(h-\bar{h})^4\rangle = -\frac{101}{105}k_{xx}(0)^4\alpha^4 t^8 - \frac{1}{6}t^5 k_{xx}(0)^2\alpha^2 k(0) + \frac{1}{2}t^2 k(0)^2, \tag{43}$$
$$\langle(h-\bar{h})^5\rangle = -(\frac{2288}{945}k_{xx}(0)^5\alpha^5 t^{10} + \frac{4}{45}k_{xx}(0)^3\alpha^3 t^7 k(0)), \tag{44}$$
$$\langle(h-\bar{h})^6\rangle = -\frac{1}{10395}t^3(85783k_{xx}(0)^6\alpha^6 t^9 + 299970k_{xx}(0)^4\alpha^4 t^6 k(0)$$
$$+623700k_{xx}(0)^2\alpha^2 t^3 k(0)^2 - 1247400k(0)^3). \tag{45}$$

The important content of the exact form derived above is that through them the time scale of sharp valley formation can be found. Actually there is no guarantee that the following generating function can be derived from a physical probability density function. So one should first check the realisability condition, i.e. $P(h-\bar{h},t) > 0$. In fact the above moment relations indicate that different even order moments become *negative* in some distinct characteristic time scales. Closer looking in the even moment relations reveals that the higher the moments are, the smaller their characteristic time scales become. So asymptotically the rate of decrease tends to $\frac{1}{4}t^*$ for very large even moments, where $t^* = (\frac{k(0)}{\alpha^2 k_{xx}^2(0)})^{1/3}$ (see figure 1). Therefore we conclude that after this time the far tails of the probability distribution function start to become negative, which is reminiscent of sharp valley creation. It means that after the characteristic time scale t^* one should also consider the contribution of the relaxation term in the limit of vanishing diffusion in order to find a realisable probability density function of height field. In other words disregarding the diffusion term in the PDF equation is valid only up to the time scales in which the singularities are developed. Taking into account that $\alpha > 0$, the odd order moments are positive in time scales before formation of sharp valleys. It means that the probability density $P(h - \bar{h}, t)$ in this time regime is negatively skewed. In figures [3-5] we have demonstrated the role of σ on the time scale of creation of singularities. Substituting the $k_{xx}(0)$ and $k(0)$ in the expression of t^* it gives us $t^* = (\pi)^{1/6} {D_0}^{-1} \alpha^{-2/3} \sigma^{5/3}$. Hence the smaller the σ is the shorter the time scale of shock creation would be(see figures 3,4 and 5).

IV. THE EQUATION OF JOINT PDF OF HEIGHT DIFFERENCE AND HEIGHT GRADIENT IN THE STATIONARY STATE

Assuming a stationary state, we are interested in investigating the stationary solutions of eq.(20) in the limit $\nu \to 0$. Of course in the stationary state the sharp valleys are fully developed and one should also take care of the diffusion term in the PDF equation. The complicated term involved with the singularities , can be cured by using the method proposed in [73]. Let us define,

$$G(u, \tilde{h}, t) = \lim_{\nu \to 0} \nu \langle u_{xx} | u, \tilde{h} \rangle P(u, \tilde{h}, t)$$

$$= \lim_{\nu \to 0} \nu \langle u_{xx}(x,t) \delta(\tilde{h} - \tilde{h}(x,t)) \delta(u - u(x,t)) \rangle, \qquad (46)$$

,where in the last step in eq.(46) we have used the definition of joint PDF $P(u, \tilde{h}, t)$. Assuming spatial ergodicity, the average of the dissipative term can be expressed as,

$$\nu\langle u_{xx}|u,\tilde{h}\rangle P(u,\tilde{h},t) = \nu\langle u_{xx}(x,t)\delta(u-u(x,t))\delta(\tilde{h}-\tilde{h}(x,t)\rangle$$
$$= \nu \lim_{L\to\infty}\frac{N}{L}\frac{1}{N}\int_{-L/2}^{L/2} dx u_{xx}(x,t)\delta(u-u(x,t))\delta(\tilde{h}-\tilde{h}(x,t)). \tag{47}$$

It is well-known that the u-field which satisfies the Burgers equation, gives rise to discontinuous or shock solutions in the limit $\nu \to 0$. Consequently for finite σ the shock solutions are manifested in height field as a set of sharp valleys at the positions where the shocks are located, where they are continuously connected by some hill configurations Fig(5). It is noted that u_{xx} is zero at the positions where no sharp valley exists. Therefore in the limit $\nu \to 0$ only small intervals around the sharp valleys will contribute to the integral in the eq.(47). Within these intervals, boundary layer analysis can be used for obtaining accurate approximation of $u(x,t), \tilde{h}(x,t)$. Generally boundary layer analysis deals with the problems in which perturbations are operative over very narrow regions across which the dependent variables undergo very rapid changes. These narrow regions (sharp valley layers) frequently adjoin the boundaries of the domain of interest, owing the fact that a small parameter (ν in the present problem) multiplies the highest derivative. A powerful method for treating boundary layer problems is the method of matched asymptotic expansions. The basic idea underlying this method is that an approximate solution to a given problem is sought not as a single expansion in terms of a single scale, but as two or more separate expansions in terms of two or more scales each of which is valid in part of the domain. The scales are chosen, so that the expansion as a whole, covers the whole domain of interest and the domains of validity of neighbouring expansions overlap. In order to handle the rapid variations in the sharp valley layers, we define a suitable magnified or stretched scale and expand the functions in terms of it in the sharp valley regions. For this purpose, we split u and h into a sum of inner solution near the sharp valleys and an outer solution away from the sharp valleys, and use systematic matched asymptotics to construct uniform approximation of u and $\tilde{h}$. For the outer solution, we look for an approximation in the form of a series in ν,

$$u = u^{out} = u_0 + \nu u_1 + O(\nu^2),$$

$$\tilde{h}= \tilde{h}^{out} = \tilde{h}_0 + \nu \tilde{h}_1 + O(\nu^2), \tag{48}$$

where u_0 and $\tilde{h}_0$ satisfy the Burgers and KPZ equation without dissipation term,

$$u_{0t} + \alpha u_0 u_{0x} = -f_x,$$

$$\tilde{h}_{0t} - \frac{\alpha}{2}(\partial_x \tilde{h}_0)^2 = f. \tag{49}$$

In order to deal with the inner solution around the sharp valley , let $y = y(t)$ be the position of a sharp valley, define the stretched variable $z = \frac{x-y}{\nu}$ and let,

$$u^{in}(x,t) = v(\frac{x-y}{\nu} + \delta, t),$$

$$\tilde{h}^{in}(x,t) = \tilde{h}(\frac{x-y}{\nu} + \delta, t). \tag{50}$$

The parameter δ is a perturbation of the sharp valley position and v and $\tilde{h}$ satisfy the following equations,

$$\nu v_t - \alpha(\bar{u} - \nu\eta) v_z + \alpha v v_z = v_{zz} - f_z(z,t),$$

$$\nu^2 \tilde{h}_t - \nu \bar{u} \tilde{h}_z + \eta \nu^2 \tilde{h}_z - \frac{\alpha}{2}(\tilde{h}_z)^2 = \nu \tilde{h}_{zz} + \nu^2 f(z,t). \tag{51}$$

$\bar{u} = \frac{1}{\alpha}\frac{dy}{dt} = \frac{u_+ + u_-}{2}, \eta = \frac{1}{\alpha}\frac{d\delta}{dt}$ and u_+, u_- are the height gradients in right hand and left hand sides of the sharp valley in the position y, (see figure-2). We look for a solution in the form,

$$v = v_0 + \nu v_1 + O(\nu^2),$$

$$\tilde{h} = \tilde{h}_0 + \nu \tilde{h}_1 + O(\nu^2). \tag{52}$$

To leading order we get for v_0 and $\tilde{h}_0$,

$$\tilde{h}_{0z} = 0,$$

$$\alpha(v_0 - \bar{u})v_{0z} = v_{0zz}, \tag{53}$$

where we have assumed that variance of $f(z,t)$ is a smooth function so that we can neglect its variation in the sharp valley region ($f_z = 0$). In other words we suppose that $\sigma >> O(\nu)$($i.e.\sigma >>$ the typical layer width). One can easily integrate the eq.(53) and find that,

$$\tilde{h}_0 = \text{const},$$
$$v_0 = \bar{u} - \frac{s}{2}\tanh(\frac{\alpha s z}{4}),$$

in which $s = s(t) = u_+ - u_-$ is the shock strength. The boundary condition for this equation arises from the matching condition,

$$\lim_{z\to\pm\infty} v_0^{in} = \lim_{x\to y} u_0^{out} = \bar{u}\pm\frac{s}{2}. \tag{54}$$

Basically since $\tilde{h}_0^{in}(z) = C - \nu\int^z v_0^{in}(z')dz'$, where C is the integration constant. So the O(1) solutions of v_0^{in} give rise to O(ν) solutions in $\tilde{h}_0^{in}$ field and only the integration constant is the O(1) part of the solution of $\tilde{h}_0^{in}$. In fact the constant in nothing but the height value at the sharp valley position. Of course due to the height continuity at sharp valley position there is no boundary layer for KPZ equation meaning that the rapid changing term in the sharp valley layer occurs in h_{xxx} while the highest derivative in KPZ equation involves only h_{xx}. The above analysis show that, to $O(\nu)$, eq.(47) can be estimated as,

$$\lim_{\nu\to 0}\nu(\langle u_{xx}|u,\tilde{h}\rangle P(u,\tilde{h},t)),$$

$$= \lim_{\nu\to 0}\nu\lim_{L\to\infty}\frac{N}{L}\frac{1}{N}\sum_j\int_{\Omega_j} dx u_{xx}^{in}\delta(u - u^{in}(x,t))\delta(\tilde{h} - \tilde{h}^{in}(y_j,t)),$$

$$= \lim_{L\to\infty}\frac{N}{L}\frac{1}{N}\sum_j\int_{-\infty}^{\infty} dz u_{zz}^{in}\delta(u - u^{in}(z,t))\delta(\tilde{h} - \tilde{h}^{in}(y_j,t)),$$

$$= \lim_{L\to\infty}\frac{N}{L}\frac{1}{N}\sum_j\int_{-\infty}^{\infty} dz v_{0zz}^{in}\delta(u - v_0)\delta(\tilde{h} - \tilde{h}(y_j,t)), \tag{55}$$

where Ω_j is a layer located at y_j with width $>> O(\nu)$. Using the eq.(53) and,

$$dzv_{0zz} = dv_0 \frac{v_{0zz}}{v_{0z}} = \alpha dv_0(v_0 - \bar{u}), \tag{56}$$

the $z-$integral can be evaluated exactly leading to the following result,

$$\nu\langle u_{xx}|u,\tilde{h}\rangle P(u,\tilde{h},t) = \alpha \int d\bar{u} \int_{-\infty}^{0} ds \varrho(\bar{u},s,\tilde{h},x,t) \int_{\bar{u}+\frac{s}{2}}^{\bar{u}-\frac{s}{2}} dv_0(v_0-\bar{u})\delta(u-v_0). \tag{57}$$

$\varrho(\bar{u},s,\tilde{h},x,t)$ is defined such that $\varrho(\bar{u},s,\tilde{h},t)d\bar{u}dsd\tilde{h}dx$ gives the average number of vallies in $[x,x+dx)$ with $\bar{u}(y,t) \in [\bar{u},\bar{u}+d\bar{u})$, $s(y,t) \in [s,s+ds)$ and $\tilde{h}(y,t) \in [\tilde{h},\tilde{h}+d\tilde{h})$, where $y \in [x,x+dx)$ is the sharp valley location. Eq.(57) indicates that the relaxation term in the strong coupling limit can be written in terms of some quantities which are defined in singularities (vallies). Indeed we characterise a sharp valley with four quantities, its location y_j, its gradients at $y_{j_{0+}}$ (i.e. u_+), $y_{j_{0-}}$ (i.e. u_-) and its height from the $\bar{h}$ i.e. $\tilde{h}_j$. Instead of u_+ and u_- we have used the quantities $\bar{u} = \frac{u_+ + u_-}{2}$ and $s = s(t) = u_+ - u_-$. Later we will determine the time evolution equations which govern over these four quantities.

Proceeding further we note that $\varrho(\bar{u},s,\tilde{h},x,t)$ can be defined as,

$$\varrho(\bar{u},s,\tilde{h},x,t) = \langle \sum_j \delta(\bar{u}-\bar{u}(y_j,t)\delta(s-s(y_j,t))\delta(\tilde{h}-\tilde{h}(y_j,t))\delta(x-y_j)\rangle. \tag{58}$$

Due to statistical homogeneity the sharp valley's characteristics are independent of their location, so

$$\varrho(\bar{u},s,\tilde{h},x,t) = \rho S(\bar{u},s,\tilde{h},t)), \tag{59}$$

in which $\rho = \rho(t)$ is the number density of shocks and $S(\bar{u},s,\tilde{h},t)$ is the PDF of $(\bar{u}(y_0,t),s(y_0,t),\tilde{h}(y_0,t))$ conditional on y_0 being a shock location. Hence

$$\lim_{\nu\to 0} \nu\langle u_{xx}|u,\tilde{h}\rangle P(u,\tilde{h},t) = -\alpha\rho \int_{-\infty}^{0} ds \int_{u+\frac{s}{2}}^{u-\frac{s}{2}} d\bar{u}(u-\bar{u})S(\bar{u},s,\tilde{h},t). \tag{60}$$

Therefore the relaxation (dissipative) contribution in eq.(20) is written as,

$$G(u,\tilde{h},t) = -(\alpha\rho \int_{-\infty}^{0} ds \int_{u+\frac{s}{2}}^{u-\frac{s}{2}} d\bar{u}(u-\bar{u})S(\bar{u},s,\tilde{h},t))_{uu}. \tag{61}$$

So eq.(20) is rewritten in the following form,

$$-P_{ut} = -\gamma P_{\tilde{h}u} - \frac{\alpha}{2}(u^2P)_{\tilde{h}u} - \alpha(uP)_{\tilde{h}} - k(0)P_{\tilde{h}\tilde{h}u} + k_{xx}(0)P_{uuu} + G(u,\tilde{h},t). \tag{62}$$

It is interesting that the the G-term comes from the relaxation term in the KPZ equation, but its explicit expression in terms of sharp valley's characteristics is proportional to α which is the coefficient of the nonlinear term in the KPZ equation. This indicates that without the nonlinear term in the KPZ equation there is no finite contribution for the diffusion term in the PDF equation when $\nu \to 0$. Although this equation is exact for finite σ however we can not solve it since the last term is not expressed in terms of $P(u,\tilde{h},t)$. Despite the existence of unclosed G-term still we can derive interesting information the moments using the above equation. we will study comprehensively the moments of height difference and height gradient, i.e. $\langle (h-\bar{h})^n(\partial_x h)^m\rangle$ in the next section.

But before it's worth of remarking that integration over $\tilde{h}$ gives an equation for probability density function (PDF) of u recovering the results in [73],

$$R_t = -k_{xx}(0)R_{uu} + \{\rho\alpha\int_{-\infty}^{0}\int_{u+\frac{s}{2}}^{u-\frac{s}{2}} d\bar{u}(u-\bar{u})S(\bar{u},s,t)\}_u \tag{63}$$

where $R(u,t) = \int P(u,\tilde{h},t)d\tilde{h}$ and $S(\bar{u},s,t) = \int S(\bar{u},s,\tilde{h},t)d\tilde{h}$ is the PDF of $(\bar{u}(y_0,t), s(y_0,t))$, conditional on the property that y_0 is the shock (cusp) position. Because of the statistical homogeneity, y_0 is a dummy variable. We finish the section with determining the relation between the density of vallies and the noise characteristics $k(0)$ and $k_{xx}(0)$, that is,

$$k_{xx}(0) = \frac{\alpha\rho}{12}\langle s^3\rangle \tag{64}$$

indicating that the forcing variance $k_{xx}(0)$ is related to the products of density of vallies ρ, $\langle s^3\rangle$ and α. This relation has been found in [73] and its details are given in the appendix.

V. THE MOMENTS OF HEIGHT FLUCTUATION IN THE STATIONARY STATE

Our goal is investigating the scaling behaviour of moments of height difference and height gradient in the stationary state. After sharp valley formation the lateral correlations produced by nonlinear term will grow with time. Dynamic scaling exponent z characterises the self-similar growth of the lateral growth. However in the stationary state the height field width saturates in the sense that lateral correlations are in average grown up to the system size. As it was explained, after the saturation the width scales as $w_0(L, t \gg L^z) \sim L^\chi$. Having in our disposal the exact result $\chi = \frac{1}{2}$ in one dimension [89], it would be natural to define $P(h', u, t)$ as PDF of h', u, and t, where $h' = \frac{h-\bar{h}}{w_0}$ and $w_0 = L^{1/2}$. Obviously $P(h', u, t)$ is related to $P(\tilde{h}, u, t)$ as $P(h', u, t) = w_0 P(\tilde{h}, u, t)$. From eq.(62) it follows that $P(h', u, t)$ in the stationary state satisfies the following equation,

$$-\gamma L^{-1/2} P_{h'u} + -\frac{\alpha}{2} L^{-1/2} (u^2 P)_{h'u} - \alpha L^{-1/2} (uP)_{h'}$$

$$-k(0) L^{-1} P_{h'h'u} + k_{xx}(0) P_{uuu} + G(u, h', t) = 0. \tag{65}$$

From eq.(65) it follows that the moments of $\langle h'^n u^m \rangle$ satisfies the following equation in the stationary state,

$$g_{n,m} + k_{xx}(0) m(m-1)(m-2) \langle h'^n u^{m-3} \rangle + mn(n-1) k(0) L^{-1} \langle h'^{n-2} u^{m-1} \rangle$$

$$- mn\gamma L^{-1/2} \langle h'^{n-1} u^{m-1} \rangle - n(m-2) \frac{\alpha}{2} L^{-1/2} \langle h'^{n-1} u^{m+1} \rangle = 0, \tag{66}$$

where

$$g_{n,m} = \int dh' \int du h'^n u^m G(u, h', t). \tag{67}$$

Using the eq.(61), we can determine the explicit expression of $g_{n,m}$ in terms of the characteristics of vallies. Thus we find,

$$g_{n,m} = -\alpha\rho \int dh' \int du h'^n u^m \Big(\int_{-\infty}^{0} ds \int_{u+\frac{s}{2}}^{u-\frac{s}{2}} d\bar{u}(u-\bar{u}) S(\bar{u}, s, h', t)\Big)_{uu}. \tag{68}$$

After integrating by parts it converts to,

$$g_{n,m} = -\alpha\rho \int dh' \int du h'^n m(m-1) u^{m-2} \Big(\int_{-\infty}^{0} ds \int_{u+\frac{s}{2}}^{u-\frac{s}{2}} d\bar{u}(u-\bar{u}) S(\bar{u}, s, h', t)\Big),$$

$$= -\alpha\rho m(m-1) \int_{-\infty}^{0} ds \int dh' h'^n \int du u^{m-2} \Big(\int_{u+\frac{s}{2}}^{u-\frac{s}{2}} d\bar{u}(u-\bar{u}) S(\bar{u}, s, h', t)\Big). \tag{69}$$

It can be integrated over u which leads to the following expression for $g_{n,m}$ (see appendix B),

$$g_{n,m} = \frac{\alpha\rho}{2^m} \big(\langle h_v'^n (2\bar{u}+s)^{m-1} [(m-1)s - 2\bar{u})]\rangle$$

$$+ \langle h_v'^n (2\bar{u}-s)^{m-1} [(m-1)s + 2\bar{u})]\rangle\big), \tag{70}$$

where $h'_v = \frac{h_v - \bar{h}}{w}$ and h_v is the height of given sharp valley . This means that the relaxation term in the strong coupling limit can be written in terms of *only* characteristics of the vallies i.e. $\bar{u}$, s and h_v.

At statistical steady state ($t \to \infty, \langle \ldots \rangle_t = 0$), and assuming the scale independence of $g_{n,m}$'s, one can derive from eq.(66), to leading order in the limit of $L \to \infty$ the following,

$$\langle (\frac{h-\bar{h}}{w_0})^n u^{m-3} \rangle = -\frac{g_{n,m}}{k_{xx}(0) m(m-1)(m-2)}, \tag{71}$$

for $m \geq 3$. Putting $n = 0$ then for instance results in the height gradient moments behaving as [73],

$$\langle u^m \rangle = \frac{\alpha\rho}{2^{m+3} k_{xx}(0)(m+1)(m+2)(m+3)}$$

$$\{\langle (2\bar{u}-s)^{m+2} [(m+2)s + 2\bar{u})]\rangle - \langle (2\bar{u}+s)^{m+2} [2\bar{u} - (m+2)s]\rangle\}. \tag{72}$$

and for $m = 3$ we find,

$$\langle (h-\bar{h})^n \rangle = \frac{g_{n,3}}{6k_{xx}(0)} \; L^{n/2} \tag{73}$$

where using the eq.(70) we have $g_{n,3} = \frac{\alpha\rho}{2}\langle h'^n s^3\rangle$. Assuming that g_{nm}'s are scale independent, at least in leading order, then one may argue that eq.(73) builds up a relation between the n-th moments of height difference in term of the second moment w_0 in a non-intermittent way. That is the n-th order moment is scaled linearly with order n. Rationalising the assumption of $g_{n,m}$'s scale independence, one would look at the statistics of sharp valley environment and the different processes involved in the sharp valley creation and annihilation which contribute dynamically. We will postpone the argumentation to next section and instead for the rest of this section we find out the consequences of scale independency of $g_{n,m}$s. In the end of this section we will verify the requirements of positivity and normalisability of the joint $P(\tilde{h}, u, t)$ in the stationary state.

Eq.(66) also suggests that the amplitudes of height difference and height gradient moments depend strongly on the singular structures in the theory, encoded in the functions $g_{n,m}$ (i.e. eq.(70)).

At this stage we find the finite size effects on the moments of $S_{n,m} = \langle h'^n u^m\rangle$. Defining $\epsilon = 1/L^{1/2}$ as a perturbative parameter we find the structure functions $S_{n,m} = \langle h'^n u^m\rangle$ perturbatively in terms of perturbative parameter $\epsilon = \frac{1}{L^{1/2}}$ as ,

$$S_{n,m} = \langle h'^n u^m\rangle = S^{(0)}_{n,m} + \epsilon S^{(1)}_{n,m} + \epsilon^2 S^{(2)}_{n,m} + \cdots, \tag{74}$$

Invoking to eq.(74) and assuming the scale independency of g_{nm} we get,

$$S^{(0)}_{n,m} = \frac{1}{k_{xx}(0)(m+1)(m+2)(m+3)} g_{n,m+3}, \tag{75}$$

$$\begin{aligned} S^{(1)}_{n,m} =& \frac{1}{k_{xx}(0)(m+1)(m+2)(m+3)}\{\frac{\gamma n(m+3)}{k_{xx}(0)(m+3)(m+4)(m+5)} g_{n-1,m+5} \\ &+ \frac{\alpha n(m+1)}{2k_{xx}(0)(m+5)(m+6)(m+7)} g_{n-1,m+7}\}, \end{aligned}$$

and,

$$S^{(2)}_{n,m} = \frac{1}{k_{xx}(0)(m+1)(m+2)(m+3)}[\gamma n(m+3) S^{(1)}_{n-1,m+4}$$

$$\frac{\alpha}{2} n(m+1) S^{(1)}_{n-1,m+4} - k(0) n(n-1)(m+3) S^{(0)}_{n-2,m+2}], \tag{76}$$

and etc. For example the moments $\langle(h-\bar{h})^n\rangle$ behave as,

$$\langle(h-\bar{h})^n\rangle = L^{n/2}\ \{\frac{1}{3!k_{xx}(0)}g_{n,3}$$

$$+\frac{1}{L^{1/2}}(-\frac{\gamma n}{5!k_{xx}(0)^2}g_{n-1,5}-\frac{2\alpha n}{7!k_{xx}(0)^2}g_{n-1,7})$$

$$+\frac{1}{L}(\frac{k(0)n(n-1)}{5!k_{xx}(0)^2}g_{n-2,5}+\frac{\gamma^2 n(n-1)}{7!k_{xx}(0)^3}g_{n-2,7}$$

$$-\frac{11\alpha\gamma n(n-1)}{9!k_{xx}(0)^2}g_{n-2,9}-\frac{40\alpha^2 n(n-1)}{11!k_{xx}(0)^2}g_{n-2,11})+O(L^{-3/2})\} \tag{77}$$

Noting to the fact that the $g_{n-1,5}$ and $g_{n-1,7}$ are not zero, therefore we conclude that the next to leading order correction for structure functions is $O(1/L^{1/2})$. Also eq. (77) shows that the amplitude of the correction terms to moments $\langle(h-\bar{h})^n\rangle$ are related to the statistics of quantities which are defined on the singularities i.e. $g_{n,m}$. Also it shows all of the moments $\langle(h-\bar{h})^n\rangle$ (for even and odd n) exist and consequently the PDF of the $h-\bar{h}$ is not symmetric. However using the properties of the Burgers equation, it can be shown that only even moments of u are nonzero and all the odd moments vanish, hence the PDF of u is symmetric.

Eq.(72) enables us to determine the rate of growth of surface at the stationary state, i.e. $\bar{h}_t$. Using the KPZ equation one it is trivial to see that,

$$\lim_{t\to\infty}\gamma(t)=\bar{h}_t=\frac{\alpha}{2}\langle u^2\rangle+\nu\rho\langle s\rangle, \tag{78}$$

where we have used the fact that $\langle h_{xx}\rangle = -\langle u_x\rangle = -\rho\langle s\rangle$ [73]. In the limit $\nu\to 0$ the second term vanishes and,

$$\lim_{t\to\infty}\gamma(t)=\bar{h}_t=\frac{\alpha}{2}\langle u^2\rangle. \tag{79}$$

Back to the eq.(72), the $\bar{h}_t$ is written in terms of properties of singularities as,

$$\bar{h}_t=\frac{\alpha^2\rho}{2^4 5!k_{xx}(0)}\{\langle 80\bar{u}^2 s^3+4s^5\rangle\} \tag{80}$$

So in the stationary state, moments $\langle \bar{u}^2 s^3 \rangle$ and $\langle s^5 \rangle$ would determine the growth rate. In other words for a given time in the steady state if one average the moments $\langle \bar{u}^2 s^3 \rangle$ and $\langle s^5 \rangle$, which is defined only on the vallies, can predict the rate of growth of surface. This provides a simple way to determine the $\bar{h}_t$ in the stationary state. Now we prove that $P(h', u, t \to \infty, L \to \infty) = P(h', u)$ is a positive and normalisable PDF. Regarding the positivity of the PDF we note that eq.(B1) indicates that $P(h', u)$ satisfies the following equation in the limit of $L \to \infty$,

$$k_{xx}(0) P_{uuu} = (\alpha\rho \int_{-\infty}^{0} ds \int_{u+\frac{s}{2}}^{u-\frac{s}{2}} d\bar{u}(u - \bar{u}) S(\bar{u}, s, \tilde{h}, t))_{uu}. \tag{81}$$

Invoking to the method introduced in section 3, one may obtain,

$$P(h', u) = -\frac{\alpha\rho}{2k_{xx}(0)} \int_{-\infty}^{0} ds \int_{u+\frac{s}{2}}^{u-\frac{s}{2}} d\bar{u}[\frac{s^2}{4} - (u - \bar{u})^2] S_{\infty}(\bar{u}, s, h'), \tag{82}$$

where $S_{\infty}(\bar{u}, s, h')$ is the PDF of vallies with $\bar{u}$, s and h'. Therefore positivity $S_{\infty}(\bar{u}, s, h') \geq 0$ implies that $P(h', u) \geq 0$. To check the normalisability of the $P(h', u)$, consider the eq.(71) with $n = 0, m = 3$, leading to,

$$\int dh' du P(h', u, t) = \frac{1}{3! k_{xx}(0)} \; g_{0,3}, \tag{83}$$

while explicit form of $g_{0,3}$ can be found from the eq.(70). The result $g_{0,3} = \frac{\alpha\rho}{2} \langle s^3 \rangle$ cooked up with the eq.(B15) gives,

$$\int dh' du P(h', u, t \to \infty, L \to \infty) = 1 \tag{84}$$

, completing the proof of normalisability of the stationary state PDF $P(h', u)$.

VI. STATISTICS FOR THE ENVIRONMENTS OF THE SINGULARITIES

In this section we derive the PDF of quantities which characterise the sharp valleys. As is depicted in Fig.(2) and formerly is described, the evolution of the surface after the formation of singularities is determined by the dynamics of the sharp vallies and their statistical properties. In a more quantitative sense one should attempt in characterising the time evolution of h_{v_j}, $\bar{u}$ and s consequently. From there we show that to leading order of the expansion in terms of system size the $g_{n,m}$'s do not depend on the scale L. Doing so we reach to such a level of describing the dynamics of the surface growth by which one may also trace the dynamics of the singularity enviornments. That makes a logical way to construct the pathway towards examining the statistical properties of singularity functions $g_{n,m}$. The importance of such an analysis became clear in the last section where the determination of the finite size corrections to scaling in $S_{n,m} = \langle h'^n u^m \rangle$ showed to be depended on the existence or lack of scale dependencies in the singularity functions $g_{n,m}$.

Let us turn to study the statistics for the environment of the singularities in KPZ equation. Defining $\xi(x,t) = -h_{xx}(x,t)$ and let $W(h_v, \bar{u}, s, \xi_+, \xi_-, x, t)$ be the PDF of

$$\begin{aligned} &h_v(x, y_j, t) = \frac{1}{2}(h(y_j + \frac{x}{2}) + h(y_j - \frac{x}{2})), \\ &\bar{u}(x, y_j, t) = \frac{1}{2}(u(y_j + \frac{x}{2}) + u(y_j - \frac{x}{2})), \\ &s(x, y_j, t) = u(y_j + \frac{x}{2}) - u(y_j - \frac{x}{2}), \\ &\xi_\pm(x, y_j, t) = -h_{x_\pm x_\pm}(y_j \pm x_\pm, t), \end{aligned}$$

conditional on y_j being a singularity position. In this section we will find the master equation governing the evolution of $W(h_v, \bar{u}, s, \xi_+, \xi_-, x, t)$ in the limit of $\nu \to 0$. Starting from the dynamical equation,

$$h_t(z + x_\pm) - \frac{\alpha}{2} h_{x_\pm}^2(z + x_\pm) = f(z + x_\pm), \tag{85}$$

$$u_t(z + x_\pm) + \alpha u u_{x_\pm}(z + x_\pm) = -f_{x_\pm}(z + x_\pm), \tag{86}$$

$$\xi_t(z + x_\pm) + \alpha u(z + x_\pm)\xi_{x_\pm}(z + x_\pm) + \alpha^2 \xi^2(z + x_\pm) = -f_{x_\pm x_\pm}(z + x_\pm), \tag{87}$$

we define,

$$\begin{aligned} &\theta(\lambda_+, \lambda_-, \mu_+, \mu_-, \eta_+, \eta_-, x_+, x_-, z, t) = \\ &\exp(-i\lambda_+ h(z + x_+) - i\lambda_- h(z + x_-) - i\mu_+ u(z + x_+) \\ &-i\mu_- u(z + x_-) - i\eta_+ \xi(z + x_+) - i\eta_- \xi(z + x_-)), \end{aligned} \tag{88}$$

and,

$$\Theta = \sum_j \theta\delta(z - y_j), \tag{89}$$

then,

$$\rho W(h_+, h_-, u_+, u_-, \xi_+, \xi_-, x, t) =$$

$$\int \frac{d\lambda_+ d\lambda_- d\mu_+ d\mu_- d\eta_+ d\eta_-}{(2\pi)^6} e^{-i\lambda_+ h_+ - i\lambda_- h_- - i\mu_+ u_+ - i\mu_- u_- - i\eta_+ \xi_+ - i\eta_- \xi_-} \langle \Theta \rangle. \tag{90}$$

We now derive equations for $\langle \Theta \rangle$ and W. Using the equations (109-111) we obtain,

$$\begin{aligned}
\langle \Theta \rangle_t = & - i\lambda_+ \langle (\frac{\alpha}{2} u_+^2 + f_+) \sum_j \theta\delta(z - y_j) \rangle \\
& - i\lambda_- \langle (\frac{\alpha}{2} u_-^2 + f_-) \sum_j \theta\delta(z - y_j) \rangle \\
& - i\mu_+ \langle (-\alpha u_+ u_{x_+} + f_{x_+}) \sum_j \theta\delta(z - y_j) \rangle \\
& - i\mu_- \langle (-\alpha u_- u_{x_-} + f_{x_-}) \sum_j \theta\delta(z - y_j) \rangle \\
& - i\eta_+ \langle (-\alpha \xi_+^2 - \alpha u_+ \xi_{x_+} + f_{x_+ x_+}) \sum_j \theta\delta(z - y_j) \rangle \\
& - i\eta_- \langle (-\alpha \xi_-^2 - \alpha u_- \xi_{x_-} + f_{x_- x_-}) \sum_j \theta\delta(z - y_j) \rangle \\
& + \langle \sum_j (-\alpha \bar{u}(y_j, t)) \delta^1(z - y_j) \theta + \sum_k (\delta(z - y_j) \delta(t - t_k) \theta \rangle \\
& - \langle \sum_l (\delta(z - y_l) \delta(t - t_l) \theta \rangle.
\end{aligned} \tag{91}$$

$\delta^1(z) = \frac{d}{dz}\delta(z)$, the (y_k, t_k)'s are the points of singularity creations and the (y_l, t_l)'s are the points of singularity annihilation due to collisions. Assuming homogeneity and using the following identity [73],

$$\theta\delta^1(z - y_j) = (\theta\delta(z - y_j))_z - \theta_{x_+} \delta(z - y_j) - \theta_{x_-} \delta(z - y_j), \tag{92}$$

it follows that,

$$\begin{aligned}
\langle \Theta \rangle_t = & - i\lambda_+ \langle (\frac{\alpha}{2} u_+^2 + f_+) \sum_j \theta\delta(z - y_j) \rangle \\
& - i\lambda_- \langle (\frac{\alpha}{2} u_-^2 + f_-) \sum_j \theta\delta(z - y_j) \rangle
\end{aligned}$$

$$
\begin{aligned}
&- i\mu_+\langle(-\alpha u_+ u_{x_+} + f_{x_+})\sum_j \theta\delta(z-y_j)\rangle\\
&- i\mu_-\langle(-\alpha u_- u_{x_-} + f_{x_-})\sum_j \theta\delta(z-y_j)\rangle\\
&- i\eta_+\langle(-\alpha\xi_+^2 - \alpha u_+\xi_{x_+} + f_{x_+x_+})\sum_j \theta\delta(z-y_j)\rangle\\
&- i\eta_-\langle(-\alpha\xi_-^2 - \alpha u_-\xi_{x_-} + f_{x_-x_-})\langle\sum_j \bar{u}(y_j,t)(\theta_{x_+}+\theta_{x_-})\delta(z-y_j)\rangle\\
&\quad +\Sigma_1 - \Sigma_2, \qquad (93)
\end{aligned}
$$

where Σ_1 and Σ_2 account respectively for singularity creation and collision events. These are given by,

$$\Sigma_1(\lambda_+,\lambda_-,\mu_+,\mu_-,\eta_+,\eta_-,x_+,x_-,z,t) = \langle\sum_k \theta\delta(z-y_k)\delta(t-t_k)\rangle, \qquad (94)$$

$$\Sigma_2(\lambda_+,\lambda_-,\mu_+,\mu_-,\eta_+,\eta_-,x_+,x_-,z,t) = \langle\sum_l \theta\delta(z-y_l)\delta(t-t_l)\rangle. \qquad (95)$$

Invoking to the Novikov's theorem we have,

$$
\begin{aligned}
\langle f_\pm\theta\rangle &= (-i\lambda_\pm k(0) - i\lambda_\mp k(x_\pm - x_\mp) - i\mu_\mp k_x(x_\pm - x_\mp)\\
&\quad + i\eta_\pm k_{xx}(0) + i\eta_\mp(x_\pm - x_\mp))\langle\theta\rangle, \qquad (96)\\
\langle f_{x_\pm}\theta\rangle &= (-i\lambda_\pm k_x(x_\pm - x_\mp) - i\mu_\pm k_{xx}(0) + i\mu_\mp k_x(x_\pm - x_\mp)\\
&\quad + i\eta_\mp k_{xxx}(x_\pm - x_\mp))\langle\theta\rangle, \qquad (97)\\
\langle f_{x_\pm x_\pm}\theta\rangle &= (-i\lambda_\pm k_{xx}(x_\pm - x_\mp)\\
&\quad - i\lambda_\mp k_{xx}(x_\pm - x_\mp) - i\mu_\mp k_{xxx}(x_\pm - x_\mp) + i\eta_\pm k_{xxxx}(0)\\
&\quad + i\eta_\mp k_{xxxx}(x_\pm - x_\mp))\langle\theta\rangle. \qquad (98)
\end{aligned}
$$

To average the convective terms we use

$$
\begin{aligned}
&i\alpha\mu_\pm\langle u_\pm\xi_\pm\Theta\rangle + i\alpha\eta_\pm\langle u_\pm\xi_\pm\Theta\rangle = -\alpha\langle u_\pm\Theta_{x_\pm}\rangle + i\alpha\lambda_\pm\langle u_\pm^2\Theta\rangle\\
&= -\alpha\langle u_\pm\Theta\rangle_{x_\pm} + \alpha\langle\xi_\pm\Theta\rangle + i\alpha\lambda_\pm\langle u_\pm^2\Theta\rangle\\
&= -i\alpha\langle\Theta\rangle_{x_\pm\mu_\pm} + i\alpha\langle\Theta\rangle_{\eta_\pm} - i\alpha\lambda_\pm\langle\Theta\rangle_{\mu_\pm\mu_\pm}. \qquad (99)
\end{aligned}
$$

Notifying that,

$$\theta_{x_\pm} = (i\lambda_\pm u_\pm - i\mu_\pm \xi_{x_\pm} - i\eta_\pm \xi_{x_\pm})\theta, \tag{100}$$

$$\langle u_\pm^2 \Theta\rangle = -\langle\Theta\rangle_{\mu_\pm\mu_\pm}, \tag{101}$$

$$\langle \xi_\pm^2 \Theta\rangle = -\langle\Theta\rangle_{\eta_\pm\eta_\pm}, \tag{102}$$

Finally for $\langle\Theta\rangle_t$ we find,

$$\begin{aligned}
\langle\Theta\rangle_t = & -\frac{i\alpha}{2}\lambda_+\langle\Theta\rangle_{\mu_+\mu_+} - \frac{i\alpha}{2}\lambda_-\langle\Theta\rangle_{\mu_-\mu_-} \\
& -\frac{i\alpha}{2}(\langle\Theta\rangle_{x_+\mu_+} + \langle\Theta\rangle_{x_-\mu_-} \\
& -\langle\Theta\rangle_{x_+\mu_-} - \langle\Theta\rangle_{x_-\mu_+} + \frac{i\alpha}{2}(\langle\Theta\rangle_{\eta_+} + \langle\Theta\rangle_{\eta_-}) \\
& -i\alpha\eta_+\langle\Theta\rangle_{\eta_+\eta_+} - i\alpha\eta_-\langle\Theta\rangle_{\eta_-\eta_-} \\
& -(\lambda_+^2 k(0) + \lambda_-^2 k(0) + 2\lambda_+\lambda_- k(x_+ - x_-))\langle\Theta\rangle \\
& -(\mu_+^2 k_{xx}(0) + \mu_-^2 k_{xx}(0) + 2\mu_+\mu_- k_{xx}(x_+ - x_-))\langle\Theta\rangle \\
& -(\eta_+^2 k_{xxxx}(0) + \eta_-^2 k_{xxxx}(0) + 2\eta_+\eta_- k(x_+ - x_-))\langle\Theta\rangle \\
& -2(\lambda_+\mu_- - \lambda_-\mu_+)k_x(x_+ - x_-)\langle\Theta\rangle + 2(\lambda_+\eta_+ + \lambda_-\eta_-)k_{xx}(0)\langle\Theta\rangle \\
& +2(\lambda_+\eta_- + \lambda_-\eta_+)k_{xx}(x_+ - x_-)\langle\Theta\rangle - 2(\eta_-\mu_+ - \eta_+\mu_-)k_{xxx}(x_+ - x_-)\langle\Theta\rangle \\
& \langle\alpha\sum_j \bar{u}(y_j,t)(\theta_{x_+} + \theta_{x_-})\delta(z - y_j)\rangle + \Sigma_1 - \Sigma_2.
\end{aligned} \tag{103}$$

for the term involving $\bar{u}(y_j,t)$ we note that,

$$u_\pm(y_j,t)\theta_{x_\pm} = (u(y_j + x_\pm,t)\theta)_{x_\pm} - \xi(y_j + x_\pm,t)\theta = i\theta_{x_\pm}\lambda_\pm - i\theta_{\mu_\pm}, \tag{104}$$

$$u_\pm(y_j,t)\theta_{x_\mp} = (u(y_j + x_\pm,t)\theta)_{x_\mp} = i\theta_{x_\mp}\lambda_\pm, \tag{105}$$

thus,

$$\begin{aligned}
&\alpha\langle\sum_j \bar{u}(y_j,t)(\theta_{x_+} + \theta_{x_-})\delta(z - y_j)\rangle \\
&= \frac{i\alpha}{2}(\langle\Theta\rangle_{x_+\lambda_+} + \langle\Theta\rangle_{x_-\lambda_+} + \langle\Theta\rangle_{x_+\lambda_-} \\
&+\langle\Theta\rangle_{x_-\lambda_-} - \langle\Theta\rangle_{\mu_+} - \langle\Theta\rangle_{\mu_-}).
\end{aligned} \tag{106}$$

Combining the above expressions, on the subset $\lambda_+ = \lambda_- = \frac{\lambda}{2}, x_+ = -x_- = \frac{x}{2}, \mu_1 = \mu_+ + \mu_-, \mu_2 = \frac{\mu_+ - \mu_-}{2}$, $\langle\Theta\rangle$ satisfies,

$$
\begin{aligned}
&\langle\Theta\rangle_t = -\frac{i\alpha}{4}\lambda(2\langle\Theta\rangle_{\mu_1\mu_1} + \frac{1}{2}\langle\Theta\rangle_{\mu_2\mu_2}) - i\alpha\langle\Theta\rangle_{x\mu_2} \\
&+\frac{i\alpha}{2}(\langle\Theta\rangle_{\eta_+} + \langle\Theta\rangle_{\eta_-}) - i\alpha\eta_+\langle\Theta\rangle_{\eta_+\eta_+} - i\alpha\eta_-\langle\Theta\rangle_{\eta_-\eta_-} \\
&-\frac{\lambda^2}{2}(k(0)+k(x))\langle\Theta\rangle - (\frac{\mu_1^2}{2} + 2\mu_2^2)k_{xx}(0) - 2(\frac{\mu_1^2}{4} - \mu_2^2)k_{xx}(x)\langle\Theta\rangle \\
&-(\eta_+^2 k_{xxxx}(0) + \eta_-^2 k_{xxxx}(0) + 2\eta_+\eta_- k_{xxxx}(x))\langle\Theta\rangle + 2\lambda\mu_2 k_x(x)\langle\Theta\rangle \\
&+\lambda(\eta_+ + \eta_-)(k_{xx}(0) + k_{xx}(x))\langle\Theta\rangle + 2\mu_2(\eta_+ + \eta_-)k_{xxx}(x)\langle\Theta\rangle + \\
&\mu_1(\eta_- - \eta_+)k_{xxx}(x)\langle\Theta\rangle + \Sigma_2 - \Sigma_1.
\end{aligned}
$$

The Σ_1, Σ_2 are evaluated at $\lambda_+ = \lambda_- = \frac{\lambda}{2}, x_+ = -x_- = \frac{x}{2}, \mu_1 = \mu_+ + \mu_-, \mu_2 = \frac{\mu_+ - \mu_-}{2}$. Changing to the variables $(h_v, \bar{u}, s, \xi_+, \xi_-)$, we obtain the following equation for W,

$$
\begin{aligned}
&(\rho W(h_v, \bar{u}, s, \xi_+, \xi_-, x, t))_t = \frac{\alpha}{2}\bar{u}^2\rho W_{h_v} + \frac{\alpha}{8}s^2\rho W_{h_v} - \alpha s\rho W_x \\
&+\frac{\alpha}{2}\xi_+\rho W + \frac{\alpha}{2}\xi_-\rho W + \alpha\rho({\xi_+}^2 W)_{\xi_+} + \alpha\rho({\xi_-}^2 W)_{\xi_-} \\
&+k(0)\rho W_{h_v h_v} + k_{xx}(0)\rho W_{\bar{u}\bar{u}} + 2\rho(k_{xx}(0) - k_{xx}(x))W_{ss} \\
&+\rho k_{xxxx}(0)(W_{\xi_+\xi_+} + W_{\xi_-\xi_-}) + 2\rho k_{xxxx}(x)W_{\xi_-\xi_-} - 2k_x(x)\rho W_{sh_v} \\
&-2k_{xx}(0)\rho W_{h_v\xi_+} - 2k_{xx}(0)\rho W_{h_v\xi_-} - 2k_{xx}(x)\rho W_{h_v\xi_+} - 2k_{xx}(x)\rho W_{h_v\xi_-} \\
&-2\rho k_{xxx}(x)(W_{s\xi_-} + W_{s\xi_+})k_{xxx}(x)\rho W_{\bar{u}\xi_-} + k_{xxx}(x)\rho W_{\bar{u}\xi_+} \\
&+\zeta_1 - \zeta_2.
\end{aligned}
\tag{107}
$$

The $\zeta_1(h_{v_j}, \bar{u}, s, \xi_+, \xi_-, x, t)$ is defined such that,

$$
\zeta_1(h_{v_j}, \bar{u}, s, \xi_+, \xi_-, x, t)dh_{v_j}dsd\bar{u}d\xi_+d\xi_-dzdt, \tag{108}
$$

gives the average number of singularity creation points in $[z, z+dz) \times [t, t+dt)$ with,

$$
\begin{aligned}
h_v(x, y_1, t_1) &\in [h_v, h_v + dh_v), \\
\bar{u}(x, y_1, t_1) &\in [\bar{u}, \bar{u} + d\bar{u}), \\
s(x, y_1, t_1) &\in [s, s + ds), \\
\xi(y_1 + \frac{x}{2}, t_1) &\in [\xi_+, \xi_+ + d\xi_+), \\
\xi(y_1 - \frac{x}{2}, t_1) &\in [\xi_-, \xi_- + d\xi_-),
\end{aligned}
$$

conditional on $(y_1,t_1) \in ([z,z+dz)\times[t,t+dt))$ being a point of singularity creation (because of the statistical homogeneity, z is a dummy variable). $\zeta_2(h_{v_j},\bar{u},s,\xi_+,\xi_-,x,t)$ is defined such that,

$$\zeta_2(h_{v_j},\bar{u},s,\xi_+,\xi_-,x,t)dh_{v_j}dsd\bar{u}d\xi_+d\xi_-dzdt, \tag{109}$$

gives the average number of singularity collision points in $[z,z+dz)\times[t,t+dt)$ with,

$$\begin{aligned} h_v(x,y_2,t_2) &\in [h_v,h_v+dh_v),\\ \bar{u}(x,y_2,t_2) &\in [\bar{u},\bar{u}+d\bar{u}),\\ s(x,y_2,t_2) &\in [s,s+ds),\\ \xi(y_2+\frac{x}{2},t_2) &\in [\xi_+,\xi_++d\xi_+),\\ \xi(y_2-\frac{x}{2},t_2) &\in [\xi_-,\xi_-+d\xi_-), \end{aligned}$$

conditional on $(y_2,t_2) \in ([z,z+dz)\times[t,t+dt))$ being a point of singularity collision. Now we rescale h_v as $h'_v = \frac{h_v}{L^{1/2}}$, so the equation (107)changes to,

$$\begin{aligned} &L^{-1/2}\frac{\alpha}{2}\bar{u}^2\rho W'_{h'_v} + L^{-1/2}\frac{\alpha}{8}s^2\rho W'_{h'_v} - \alpha s\rho W'_x + \frac{\alpha}{2}\xi_+\rho W' + \frac{\alpha}{2}\xi_-\rho W'\\ &+\alpha\rho({\xi_+}^2W')_{\xi_+} + \alpha\rho({\xi_-}^2W')_{\xi_-} + L^{-1}k(0)\rho W'_{h'_vh'_v} + k_{xx}(0)\rho W'_{\bar{u}\bar{u}}\\ &+2\rho(k_{xx}(0)-k_{xx}(x))W'_{ss} + \rho k_{xxxx}(0)(W'_{\xi_+\xi_+} + W'_{\xi_-\xi_-})\\ &+2\rho k_{xxxx}(x)W'_{\xi_-\xi_-} - 2L^{-1/2}k_x(x)\rho W'_{sh'_v} - 2k_{xx}(0)L^{-1/2}\rho W'_{h'_v\xi_+}\\ &-2k_{xx}(0)L^{-1/2}\rho W'_{h'_v\xi_-} - 2k_{xx}(x)L^{-1/2}\rho W'_{h'_v\xi_+} - 2k_{xx}(x)L^{-1/2}\rho W'_{h'_v\xi_-}\\ &-2\rho k_{xxx}(x)(W'_{s\xi_-} + W'_{s\xi_+}) - k_{xxx}(x)\rho W'_{\bar{u}\xi_-} + k_{xxx}(x)\rho W'_{\bar{u}\xi_+}\\ &+\zeta'_1 - \zeta'_2 = 0. \end{aligned} \tag{110}$$

In the limit of large L or $L\to\infty$ the leading terms are,

$$\begin{aligned} &-\alpha s\rho W'_x + \frac{\alpha}{2}\xi_+\rho W' + \frac{\alpha}{2}\xi_-\rho W'\\ &+\alpha\rho({\xi_+}^2W')_{\xi_+} + \alpha\rho({\xi_-}^2W')_{\xi_-} + k_{xx}(0)\rho W'_{\bar{u}\bar{u}}\\ &+2\rho(k_{xx}(0)-k_{xx}(x))W'_{ss} + \rho k_{xxxx}(0)(W'_{\xi_+\xi_+} + W'_{\xi_-\xi_-}) \end{aligned}$$

$$+2\rho k_{xxxx}(x) W'_{\xi_-\xi_-} - 2\rho k_{xxx}(x)(W'_{s\xi_-} + W'_{s\xi_+}) -$$

$$k_{xxx}(x)\rho W'_{\bar{u}\xi_-} + k_{xxx}(x)\rho W'_{\bar{u}\xi_+} + \zeta'_1 - \zeta'_2 = 0. \tag{111}$$

To find the $g_{n,m}$ we multiply the above equation by $h'^n \bar{u}^m s^p$ and integrating over $h', \bar{u}, s, \xi_+$ and ξ_- we have,

$$-\alpha\rho\langle h_v^{'n} \bar{u}^m s^{p+1}\rangle_x + \frac{\alpha\rho}{2}\langle h_v^{'n} \bar{u}^m s^p \xi_+\rangle + \frac{\alpha\rho}{2}\langle h_v^{'n} \bar{u}^m s^p \xi_-\rangle$$

$$+2p(p-1)\rho[k_{xx}(0) - k_{xx}(x)]\langle h_v^{'n} \bar{u}^m s^{p-2}\rangle$$

$$+m(m-1)k_{xx}(0)\rho\langle h_v^{'n} \bar{u}^{m-2} s^p\rangle + Q^{(1)}_{nmp} - Q^{(2)}_{nmp} = 0, \tag{112}$$

where,

$$Q^{(1)}_{nmp} = \int h'^n \bar{u}^m s^p \zeta'_1 dh' d\bar{u} ds d\xi_+ d\xi_-,$$

$$Q^{(2)}_{nmp} = \int h'^n \bar{u}^m s^p \zeta'_2 dh' d\bar{u} ds d\xi_+ d\xi_-.$$

Using the following identity,

$$\frac{\partial}{\partial x} h'_v(x, y_j, t) = -\frac{s}{4L^{1/2}}, \tag{113}$$

$$\frac{\partial}{\partial x} s(x, y_j, t) = \frac{1}{2}(\xi_+ + \xi_-), \tag{114}$$

$$\frac{\partial}{\partial x} \bar{u}(x, y_j, t) = \frac{s}{2}, \tag{115}$$

we find,

$$\frac{\alpha\rho}{2}\langle h_v^{'n} \bar{u}^m s^p(\xi_+ + \xi_-)\rangle = \frac{\alpha\rho}{p+1}\Big\{\langle h_v^{'n} \bar{u}^m s^{p+1}\rangle_x$$

$$+ \frac{n}{4L^{S/2}}\langle h_v^{'(n-1)} S \bar{u}^m s^{p+2}\rangle + \frac{m}{2}\langle h_v^{;n} \bar{u}^{m-1} s^{p+2}\rangle\Big\} \tag{116}$$

So in the limit of $L \rightarrow \infty$ we have

$$-\frac{p\alpha\rho}{p+1}\langle h_v'^n\bar{u}^m s^{p+1}\rangle_x + \frac{m\alpha\rho}{2(p+1)}\langle h_v'^n\bar{u}^{m-1}s^{p+2}\rangle$$

$$+2p(p-1)\rho[k_{xx}(0)-k_{xx}(x)]\langle h_v'^n\bar{u}^m s^{p-2}\rangle$$

$$+m(m-1)k_{xx}(0)\rho\langle h_v'^n\bar{u}^{m-2}s^p\rangle + Q^{(1)}_{nmp} - Q^{(2)}_{nmp} = 0. \tag{117}$$

Assuming a stationary solution for the dynamical equation (107) governed over W and rescaling h_v as $h_v' = \frac{h_v}{L^{1/2}}$ in the resulting differential equation we reach to eq.(110). Of course the mentioned equation *is* dependent on scale L but being interested in the limit of $L \to \infty$ results to eq.(111) which is free of the explicit scale dependent terms in the leading order . However we are faced with two very complicated terms namely ζ_1 and ζ_2 which would be analysed . The origin of these terms are related to processes of sharp valley creation and annihilation. We argue that these processes are basically involving local interaction between nearby sharp valleys and effects of forcing, which its spatial correlation is assumed to be much less than system size, so they essentially would not carry any information about system size. In this sense eq.(111) encodes the fact that the probability distribution W is a scale invariant function of its argument $h_v' = \frac{h_v}{L^{1/2}}$ in the leading order. The above property is deciphered in eq.(117) too but this time it is translated in terms of the scale independence of $g_{n,m}$s.

Also the equation for W enables us to find the time evolution of the sharp valley characteristics. For example multiplying the equation (107) by h_v and integrating over all variables we can derive the increasing rate of mean height of the singularities and noting that,

$$\frac{\partial}{\partial x}h_v(x,y_j,t) = -\frac{s}{4}, \tag{118}$$

$$\frac{\partial}{\partial x}s(x,y_j,t) = \frac{1}{2}(\xi_+ + \xi_-), \tag{119}$$

we get,

$$\frac{d}{dt}\langle h_v\rangle(t) = -\frac{\alpha}{8}\langle 4\bar{u}^2 - s^2\rangle. \tag{120}$$

VII. CONCLUSION

We study the problem of non-equilibrium surface growth described by forced KPZ equation in 1+1 dimensions. The forcing is a white in time Gauaaian noise but with a Gaussian correlation in space. Modelling a short range correlated noise we restrict our study to the case when the correlation length of the forcing is much smaller than the system size. In the non-stationary regime when the sharp valley structures are not yet developed we find an exact form for the generating function of the joint fluctuations of height and height gradient. We determine the time scale of the sharp valley formation and the exact functional form of the time dependence in the height difference moments at any given order. Investigating the stationary state we give a general expression of the mixed correlations of height and height-gradient at any order, in terms of the quantities which characterise the sharp valley singular structures. Through a careful analysis being done over the behaviour of the sharp valley environment, we decipher the general finite size corrections to the scaling of an arbitrary nth moment, i.e. $\langle (h-\bar{h})^n \rangle$, at any order. Recently Marinari etal. [23] have obtained the corrections to the leading order scaling in dimensions $D = 2, 3, 4$, in a high resolution simulation on the RSOS discrete model which is beleived to be in the universality class of the KPZ equation stirred with a white in time Gaussian noise and delta correlated in space. Hence they get,

$$w_n(L) \sim A_n L^{n\chi}(1 + B_n L^{-\omega}). \tag{121}$$

Irrespective of the dimension and moment order n, they observe the same sub-leading exponent ω always very close to unity (see also [91,92]). Through our calculations we succeed to obtain the finite size corrections analytically. However we have to remark that, due to working with finite correlated forcing a firm comparison between our results and numerical simulations is not possible. More precisely, in the present paper the limiting of $\nu \to 0$ is taken into account only when σ is finite. Still the forcing correlation length is much smaller than the system size and height correlation length. But the limiting of $\sigma \to 0$ is a singular limit in our calculations, and moreover, it is not a priori clear that the limits of $\nu \to 0$ and $\sigma \to 0$ commute at all. However due to the scale independence of $g_{n,m}$'s, the eq.(77) shows the general correction terms for nth order moment, all having the same sub-leading exponent $\omega = 1/2$. The amplitudes A_n and B_n in eq.(121) are given explicitly in terms of the functions $g_{n,m}$ defined on the sharp valley singularities. The next step, left for the future, would be the calculation of $g_{n,m}$'s in terms of few known parameters, i.e. the forcing and diffusion coefficients.

Our analysis enables us to find the stochastic equations which are governed over the dynamics of quantities characterising the cusp singularities too. This translates the stationary non-equilibrium dynamics of the surface in terms of the dynamics of singularities in the stationary state. When the system crosses over the time t^*, after which the first singularties are formed, it would be an important study to analyse the shape deformation of non-stationary height PDF $P(h', t)$ in time. We believe that the analysis followed in this paper is quite suitable for the zero temperature limit in the problem of directed polymer in the random potential with short range correlations [88]. The same method applied to KPZ equation in higher dimensions would be definitely one of the consequent goals of the present work. The main message which might be encoded in the present work is the importance of the statistical properties of the geometrical singular structures for understanding the strong coupling regime of Kardar-Parisi-Zhang equation in higher dimensions.

Acknowledgement

We would like to appreaciate J. Bec for kindly providing us some of his unpublished numerical results. We also thank C. Castellano for reading the manuscript and bringing our attention to some of the references and Roya Mohayaee for helpful comments. We are indebted to Eric Vanden Eijnden for very informative discussions.

APPENDIX A: AN ALTERNATIVE METHOD FOR DETERMINING OF THE MOMENTS OF HEIGHT FLUCTUATION BEFORE THE FORMATION OF THE SINGULARITIES

In this appendix we give the details of calculations of the scaling behaviour of moments of height difference before the formation of singularities. We know that the generating function $Z(\mu, \lambda, t)$ satisfies the following equation when $(\nu \to 0)$,

$$Z_t = i\gamma\lambda Z - \frac{i\lambda\alpha}{2} Z_{\mu\mu} - \lambda^2 k(0) Z + \frac{i\alpha\lambda}{\mu} Z_\mu + \mu^2 k_{xx}(0) Z. \tag{A1}$$

Let us write $Z(\mu, \lambda, t)$ as follow,

$$\begin{aligned} Z(\mu, \lambda, t) &:= (1 + A(t)\lambda^2 + C(t)\lambda^3 + F(t)\lambda^4 + G(t)\lambda^5 + J(t)\lambda^6 + M(t)\lambda^7 \\ &+ B(t)\lambda\mu^2 + D(t)\lambda^2\mu^2 + E(t)\lambda\mu^4 + H(t)\lambda^3\mu^2 + K(t)\lambda^2\mu^4 + L(t)\lambda^4\mu^2 + \\ &N(t)\lambda^5\mu^2 + P(t)\lambda^3\mu^4 + Q(t)\lambda\mu^6) \exp((-\lambda^2 k(0) + \mu^2 k_{xx}(0))t). \end{aligned} \tag{A2}$$

Now expanding $Z(\mu, \lambda, t)$ as a series of μ, λ and substituting it in the equation eq.(26), we equate the terms in different orders of μ, λ ending with some coupled differential equations governed over the coefficients introduced in the definition of Z in eq.(26). So we have,

$$\frac{\partial}{\partial t} A(t) = i\alpha B(t), \tag{A3}$$

$$\frac{\partial}{\partial t} B(t) = -2i\alpha k_{xx}(0)^2 t^2, \tag{A4}$$

$$\frac{\partial}{\partial t} C(t) = i\alpha D(t), \tag{A5}$$

$$\frac{\partial}{\partial t} D(t) = -4i\alpha k_{xx}(0) t B(t) - 2i\alpha E(t), \tag{A6}$$

$$\frac{\partial}{\partial t} E(t) = 0, \tag{A7}$$

$$\frac{\partial}{\partial t} F(t) = i\alpha H(t), \tag{A8}$$

$$\frac{\partial}{\partial t} H(t) = -4i\alpha k_{xx}(0) t D(t) - 2i\alpha K(t) - 2i\alpha k_{xx}(0)^2 t^2 A(t), \tag{A9}$$

$$\frac{\partial}{\partial t}K(t) = -2i\alpha k_{xx}(0)^2 t^2 B(t) - 8i\alpha k_{xx}(0) t E(t) - 9i\alpha Q(t), \tag{A10}$$

$$\frac{\partial}{\partial t}Q(t) = 0, \tag{A11}$$

By solving these differential equations with the initial conditions that $A(t), B(t), C(t), D(t), E(t), F(t), H(t), K(t), Q(t)$ are zero at $t = 0$, we find,

$$A(t) = \frac{1}{6}\alpha^2 k_{xx}(0)^2 t^4, \tag{A12}$$

$$B(t) = -\frac{2}{3}i\alpha k_{xx}(0)^2 t^3, \tag{A13}$$

$$C(t) = \frac{4}{45}i\alpha^3 k_{xx}(0)^3 t^6, \tag{A14}$$

$$D(t) = \frac{8}{15}\alpha^2 k_{xx}(0)^3 t^5, \tag{A15}$$

$$E(t) = 0, \tag{A16}$$

$$F(t) = -\frac{101}{2520}\alpha^4 k_{xx}^4 t^8, \tag{A17}$$

$$H(t) = \frac{101}{315}i\alpha^3 k_{xx}^4 t^7, \tag{A18}$$

$$K(t) = -\frac{2}{9}\alpha^2 k_{xx}(0)^4 t^6, \tag{A19}$$

$$Q(t) = 0. \tag{A20}$$

By replacing these expressions in eq.(A2) we find $Z(\mu, \lambda, t)$ as a function of μ, λ, t explicitly without any unknown terms or expressions. Now if we expand the original form of generating function $Z(\mu, \lambda, t)$ as a series in μ, λ we find,

$$\begin{aligned}
&Z(\mu, \lambda, t) = \langle \exp i\lambda((h - \bar{h})) + i\mu(\partial_x(h - \bar{h}))\rangle = \\
&-\frac{1}{720}u^6\mu^6 - \frac{1}{120}(h - \bar{h})u^5\mu^5\lambda - \frac{1}{48}(h - \bar{h})^2 u^4\mu^4\lambda^2 - \frac{1}{36}(h - \bar{h})^3 u^3\mu^3\lambda^3 \\
&-\frac{1}{48}(h - \bar{h})^4 u^2\mu^2\lambda^4 - \frac{1}{120}(h - \bar{h})^5 u\mu\lambda^5 - \frac{1}{720}(h - \bar{h})^6\lambda^6 - \frac{1}{120}iu^5\mu^5 \\
&-\frac{1}{24}i(h - \bar{h})u^4\mu^4\lambda - \frac{1}{12}i(h - \bar{h})^2 u^3\mu^3\lambda^2 - \frac{1}{12}i(h - \bar{h})^3 u^2\mu^2\lambda^3 - \frac{1}{24}i(h - \bar{h})^4 u\mu\lambda^4 \\
&-\frac{1}{120}i(h - \bar{h})^5\lambda^5 + \frac{1}{24}u^4\mu^4 + \frac{1}{6}(h - \bar{h})u^3\mu^3\lambda + \frac{1}{4}(h - \bar{h})^2 u^2\mu^2\lambda^2 \\
&+\frac{1}{6}(h - \bar{h})^3 u\mu\lambda^3 + \frac{1}{24}(h - \bar{h})^4\lambda^4 + \frac{1}{6}iu^3\mu^3 + \frac{1}{2}i(h - \bar{h})u^2\mu^2\lambda
\end{aligned}$$

$$+\frac{1}{2}i(h-\bar{h})^2u\mu\lambda^2+\frac{1}{6}i(h-\bar{h})^3\lambda^3-\frac{1}{2}u^2\mu^2-(h-\bar{h})u\mu\lambda$$
$$-\frac{1}{2}(h-\bar{h})^2\lambda^2-iu\mu-i(h-\bar{h})\lambda+1. \tag{A21}$$

Equating the coefficients of eq.(A2) and eq.(A21) proportional to the same powers in μ and λ and replacing the expressions of $A(t), B(t), C(t), D(t), E(t), F(t), H(t), K(t), Q(t)$ it results to the same expressions given before, i.e.

$$\langle(h-\bar{h})^2\rangle = -\frac{1}{3}t(k_{xx}(0)^2\alpha^2t^3-6k(0)), \tag{A22}$$
$$\langle(h-\bar{h})^3\rangle = -\frac{24}{45}k_{xx}(0)^3\alpha^3t^6, \tag{A23}$$
$$\langle(h-\bar{h})^4\rangle = -\frac{101}{105}k_{xx}(0)^4\alpha^4t^8-\frac{1}{6}t^5k_{xx}(0)^2\alpha^2k(0)+\frac{1}{2}t^2k(0)^2. \tag{A24}$$

APPENDIX B: THE PROOF OF RELATION BETWEEN ρ AND $\langle S^3\rangle$

We consider the statistical steady state i.e. $R_t=0$ so that the eq.(63) can be written as follows,

$$R_{uu}=\frac{1}{k_{xx}(0)}G(u,t)=\frac{\alpha\rho}{k_{xx}(0)}(\int_{-\infty}^{0}ds\int_{u+\frac{s}{2}}^{u-\frac{s}{2}}d\bar{u}(u-\bar{u})S(\bar{u},s,t))_u. \tag{B1}$$

We integrate the eq.(B1) w.r.t u and find,

$$R_u=\frac{\alpha\rho}{k_{xx}(0)}\int_{-\infty}^{0}ds\int_{u+\frac{s}{2}}^{u-\frac{s}{2}}d\bar{u}(u-\bar{u})S(\bar{u},s,t). \tag{B2}$$

At the large time limit ($t\to\infty$) we denote R and S as R_∞ and $S_\infty(\bar{u},s)$. So,

$$R_\infty=\frac{\alpha\rho}{k_{xx}(0)}\int_{-\infty}^{u}du\int_{-\infty}^{0}ds\int_{u+\frac{s}{2}}^{u-\frac{s}{2}}d\bar{u}(u-\bar{u})S_\infty(\bar{u},s). \tag{B3}$$

To determine the R_∞ we define function $K(u)$ as the following,

$$K(u)=\frac{\alpha\rho}{k_{xx}(0)}\int_{-\infty}^{0}ds\int_{u+\frac{s}{2}}^{u-\frac{s}{2}}d\bar{u}\frac{(u-\bar{u})^2}{2}S_\infty(\bar{u},s). \tag{B4}$$

Differentiation the above equation with respect to u gives us,

$$\frac{d}{du}K(u)=\frac{\alpha\rho}{k_{xx}(0)}\int_{-\infty}^{0}ds\int_{u+\frac{s}{2}}^{u-\frac{s}{2}}d\bar{u}(u-\bar{u})S_\infty(\bar{u},s)$$

$$\frac{\alpha\rho}{k_{xx}(0)}\int_{-\infty}^{0}ds\frac{s^2}{8}S_\infty(u-\frac{s}{2},s)-\frac{\alpha\rho}{k_{xx}(0)}\int_{-\infty}^{0}ds\frac{s^2}{8}S_\infty(u+\frac{s}{2},s). \tag{B5}$$

Now we integrate the eq.(B5) over u from $-\infty$ to u and find,

$$\int_{-\infty}^{u}du\frac{d}{du}K(u)=\frac{\alpha\rho}{k_{xx}(0)}\int_{-\infty}^{u}du\int_{-\infty}^{0}ds\int_{u+\frac{s}{2}}^{u-\frac{s}{2}}d\bar{u}(u-\bar{u})S_\infty(\bar{u},s)$$

$$\frac{\alpha\rho}{2k_{xx}(0)}\int_{-\infty}^{u}du\int_{-\infty}^{0}ds\frac{s^2}{4}S_\infty(u-\frac{s}{2},s)-\frac{\alpha\rho}{2k_{xx}(0)}\int_{-\infty}^{u}du\int_{-\infty}^{0}ds\frac{s^2}{4}S_\infty(u+\frac{s}{2},s). \tag{B6}$$

Then we will find,

$$K(u)-K(-\infty)=R_\infty(u)+\frac{\alpha\rho}{2k_{xx}(0)}\int_{-\infty}^{0}ds\int_{-\infty}^{u-\frac{s}{2}}d\bar{u}\frac{s^2}{4}S_\infty(\bar{u},s)$$

$$-\frac{\alpha\rho}{2k_{xx}(0)}\int_{-\infty}^{0}ds\int_{-\infty}^{u+\frac{s}{2}}d\bar{u}\frac{s^2}{4}S_\infty(\bar{u},s). \tag{B7}$$

According to the definition of $K(u)$ we see that $K(-\infty)\to 0$ (the shock probability density function goes to zero in this limit) and therefore we find the following relation between $K(u)$ and $R_\infty(u)$,

$$K(u)=R_\infty(u)+\frac{\alpha\rho}{2k_{xx}(0)}\int_{-\infty}^{0}ds\int_{u+\frac{s}{2}}^{u-\frac{s}{2}}d\bar{u}\frac{s^2}{4}S_\infty(\bar{u},s). \tag{B8}$$

Using the eqs.(B4) and (B8) we find explicit relation between the R_∞ and $S_\infty(\bar{u},s)$ as follows,

$$R_\infty(u)=-\frac{\alpha\rho}{2k_{xx}(0)}\int_{-\infty}^{0}ds\int_{u+\frac{s}{2}}^{u-\frac{s}{2}}d\bar{u}[\frac{s^2}{4}-(u-\bar{u})^2]S_\infty(\bar{u},s). \tag{B9}$$

Assuming $S_\infty(\bar{u},s)\geq 0$, it becomes evident that above integral would give a realisable portability density for height gradient, that is $R_\infty\geq 0$. For finite σ the eq.(B9) gives us the PDF of height gradient in the KPZ equation in the strong coupling limit. The function $R_\infty(u)$) enables us to determine the relation between the vallies density ρ and $k_{xx}(0)$. We would integrate over u from R_∞ so we define another function $K_1(u)$ such that,

$$K_1(u)=-\frac{\alpha\rho}{2k_{xx}(0)}\int_{-\infty}^{0}ds\int_{u+\frac{s}{2}}^{u-\frac{s}{2}}d\bar{u}[\frac{s^2}{4}u-\frac{(u-\bar{u})^3}{3}]S_\infty(\bar{u},s), \tag{B10}$$

where differentiation $K_1(u)$ with respect to u gives,

$$\frac{d}{du}K_1(u) = -\frac{\alpha\rho}{2k_{xx}(0)}\int_{-\infty}^{0} ds \int_{u+\frac{s}{2}}^{u-\frac{s}{2}} d\bar{u}[\frac{s^2}{4} - (u-\bar{u})^2]S_\infty(\bar{u}, s)$$

$$-\frac{\alpha\rho}{2k_{xx}(0)}\int_{-\infty}^{0} ds[\frac{s^2}{4}u - \frac{s^3}{24}]S_\infty(u - \frac{s}{2}, s)$$

$$+\frac{\alpha\rho}{2k_{xx}(0)}\int_{-\infty}^{0} ds[\frac{s^2}{4}u + \frac{s^3}{24}]S_\infty(u + \frac{s}{2}, s). \tag{B11}$$

Now integrating the above equation over u from $-\infty$ to $+\infty$ gives,

$$K_1(+\infty) - K_1(-\infty) = \int_{-\infty}^{+\infty} du R_\infty(u)$$

$$-\frac{\alpha\rho}{2k_{xx}(0)}\int_{-\infty}^{+\infty} du \int_{-\infty}^{0} ds[\frac{s^2}{4}(u + \frac{s}{2}) - \frac{s^3}{24}]S_\infty(u, s)$$

$$+\frac{\alpha\rho}{2k_{xx}(0)}\int_{-\infty}^{+\infty} du \int_{-\infty}^{0} ds[\frac{s^2}{4}(u - \frac{s}{2}) + \frac{s^3}{24}]S_\infty(u, s). \tag{B12}$$

Using the fact that $K_1(+\infty) = K_1(-\infty) = 0$, we obtain,

$$\int_{-\infty}^{+\infty} du R_\infty(u) = -\frac{\alpha\rho}{2k_{xx}(0)}\int_{-\infty}^{+\infty} du \int_{-\infty}^{0} ds[\frac{s^2}{4}u - \frac{s^3}{12}]S_\infty(u, s)$$

$$+\frac{\alpha\rho}{2k_{xx}(0)}\int_{-\infty}^{+\infty} du \int_{-\infty}^{0} ds[\frac{s^2}{4}u + \frac{s^3}{12}]S_\infty(u, s), \tag{B13}$$

in which the sum of the terms in right hand side gives,

$$\int_{-\infty}^{+\infty} du R_\infty(u) = \frac{\alpha\rho}{12k_{xx}(0)}\langle s^3\rangle. \tag{B14}$$

Thus from the requirement that R_∞ be normalised to unity, we get,

$$k_{xx}(0) = \frac{\alpha\rho}{12}\langle s^3\rangle. \tag{B15}$$

APPENDIX C: DERIVATION OF THE FINITE CONTRIBUTION OF RELAXATION TERM IN THE STATIONARY STATE

In this appendix we give the detail of calculations of $g_{n,m}$ in the equation eq.(70). For computing the g_{nm} we introduce,

$$K(u) = -m(m-1)\alpha\rho \int_{-\infty}^{0} ds \int_{-\infty}^{\infty} dh' h'^{n} \int_{u-\frac{s}{2}}^{u+\frac{s}{2}} d\bar{u} \{ \frac{u^{m-1}}{m-1}\bar{u} - \frac{u^{m}}{m} \} S(\bar{u}, s, h', t). \tag{C1}$$

By differentiating $K(u)$ and integrating in the whole range of u we have,

$$\int_{-\infty}^{\infty} \frac{dK(u)}{du} du = g_{nm}$$

$$-m(m-1)\alpha\rho \int_{-\infty}^{0} ds \int_{-\infty}^{\infty} dh' h'^{n} \int_{-\infty}^{\infty} du \{ \frac{u^{m}}{m} - \frac{u^{m-1}}{m-1}(u - \frac{s}{2}) \} S(u - \frac{s}{2}, s, h', t)$$

$$-m(m-1)\alpha\rho \int_{-\infty}^{0} ds \int_{-\infty}^{\infty} dh' h'^{n} \int_{-\infty}^{\infty} du \{ \frac{u^{m}}{m} - \frac{u^{m-1}}{m-1}(u + \frac{s}{2}) \} S(u + \frac{s}{2}, s, h', t). \tag{C2}$$

Since $K(+\infty) = K(-\infty) = 0$, left hand side vanishes, so,

$$g_{nm} = m(m-1)\alpha\rho \int_{-\infty}^{0} ds \int_{-\infty}^{\infty} dh' h'^{n} \int_{-\infty}^{\infty} d\bar{u} [\frac{(\bar{u} + \frac{s}{2})^{m}}{m} - \frac{(\bar{u} + \frac{s}{2})^{m-1}}{m-1} \bar{u}] S(\bar{u}, s, h', t)$$

$$- m(m-1)\alpha\rho \int_{-\infty}^{0} ds \int_{-\infty}^{\infty} dh' h'^{n} \int_{-\infty}^{\infty} d\bar{u} [\frac{(\bar{u} - \frac{s}{2})^{m}}{m} - \frac{(\bar{u} - \frac{s}{2})^{m-1}}{m-1} \bar{u}] S(\bar{u}, s, h', t)$$

$$= \frac{\rho}{2^{m}} \int_{-\infty}^{0} ds \int_{-\infty}^{\infty} d\bar{u} \int_{-\infty}^{\infty} dh' h'^{n} \{ (2\bar{u} + s)^{m-1}$$

$$\{ [(m-1)s - 2\bar{u}] + (2\bar{u} - s)^{m-1} [(m-1)s + 2\bar{u}] \} S(\bar{u}, s, h', t), \tag{C3}$$

which finally leads to eq.(70).

APPENDIX D: DYNAMICS OF QUANTITIES WHICH ARE DEFINED ON THE SHARP VALLEYS

In this section we determine the equation of motion for $\bar{u}(y_j, t), s(y_j, t), \tilde{h}(y_j, t)$ along the sharp valley which is located at position y_j at time t. Using the KPZ equation and its differentiation by x around the sharp valley at position y_j, one can find a set of equation for $h_+(y_j,t) = \lim_{\epsilon\to 0^+} h(y_j+\epsilon,t)$, $h_-(y_j,t) = \lim_{\epsilon\to 0^+} h(y_j-\epsilon,t)$, $u_+(y_j,t) = \lim_{\epsilon\to 0^+} u(y_j+\epsilon,t)$, and $u_-(y_j,t) = \lim_{\epsilon\to 0^+} u(y_j-\epsilon,t)$ as follows,

$$h_{+t}(y_j,t) = \frac{\alpha}{2}u_+^2(y_j,t) + f(y_j,t), \tag{D1}$$

$$h_{-t}(y_j,t) = \frac{\alpha}{2}u_-^2(y_j,t) + f(y_j,t), \tag{D2}$$

$$u_{+t}(y_j,t) = -\alpha u_+(y_j,t)u_{+x}(y_j,t) - f_x(y_j,t), \tag{D3}$$

$$u_{-t}(y_j,t) = -\alpha u_-(y_j,t)u_{-x}(y_j,t) - f_x(y_j,t). \tag{D4}$$

To determine the $\frac{d}{dt}\{\bar{u}, s, \tilde{h}\}$ we use the following identity [73,84],

$$\begin{aligned}\frac{d}{dt}u_+(y_j,t) &= \frac{dy_j}{dt}u_{+x}(y_j,t) + u_{+t}(y_j,t)\\ &= \alpha\bar{u}(y_j,t)u_{+x}(y_j,t) - \alpha u_{+x}(y_j,t)u_+(y_j,t) - f_x(y_j,t)\\ &= -\frac{\alpha}{2}s(y_j,t)u_{+x}(y_j,t) - f_x(y_j,t),\end{aligned} \tag{D5}$$

where $\bar{u} = \frac{1}{\alpha}\frac{dy_j}{dt}$. Similarly,

$$\frac{d}{dt}\bar{u}(y_j,t) = \frac{\alpha}{2}s(y_j,t)u_{-x}(y_j,t) - f_x(y_j,t). \tag{D6}$$

These equations can be re-written as,

$$\frac{d}{dt}\bar{u}(y_j,t) = -\frac{\alpha}{4}s(u_{+x} - u_{-x}) - f_x,$$

$$\frac{d}{dt}s(y_j,t) = -\frac{\alpha}{2}s(u_{+x} + u_{-x}), \tag{D7}$$

where will give the equations for $\bar{u}$ and s. Since $u = -h_x$ we write the above equations in term of curvature of the surface in the right and left sides of the sharp valley at position y_j as,

$$\frac{d}{dt}\bar{u}(y_j,t) = \frac{\alpha}{4}s(h_{+xx} - h_{-xx}) - f_x,$$

$$\frac{d}{dt}s(y_j,t) = \frac{\alpha}{2}s(h_{+xx} + h_{-xx}). \tag{D8}$$

For determining the time evolution of the $\tilde{h} = h - \bar{h}$, we use the KPZ equation by which one can easily show that $h_+(y_j, t)$ and $h_-(y_j, t)$ satisfy,

$$\frac{d}{dt}h_+(y_j,t) = \frac{dy_j}{dt}h_{+x}(y_j,t) + h_{+t}(y_j,t),$$

$$\frac{d}{dt}h_-(y_j,t) = \frac{dy_j}{dt}h_{-x}(y_j,t) + h_{-t}(y_j,t). \tag{D9}$$

By definition we have $\frac{d}{dt}h(y_j,t) = \frac{1}{2}(\frac{d}{dt}h_+ + \frac{d}{dt}h_-)$, so using the equation for h_+ and h_-,

$$\frac{d}{dt}h(y_j,t) = -\frac{\alpha}{8}(4\bar{u}^2 - s^2) + f, \tag{D10}$$

and

$$\frac{d}{dt}\tilde{h}(y_j,t) = -\frac{\alpha}{8}(4\bar{u}^2 - s^2) + f - \gamma, \tag{D11}$$

where $\tilde{h}(y_j,t) = h(y_j,t) - \bar{h}$ and $\bar{h}_t = \gamma$.

Therefore in summary we have the following set of equations for given sharp valley in the KPZ problem in the limit $\nu \rightarrow 0$,

$$\frac{dy_j}{dt} = \alpha\bar{u},$$

$$\frac{d}{dt}\bar{u}(y_j,t) = \frac{\alpha}{4}s(h_{+xx} - h_{-xx}) - f_x,$$

$$\frac{d}{dt}s(y_j,t) = \frac{\alpha}{2}s(h_{+xx} + h_{-xx}),$$

$$\frac{d}{dt}\tilde{h}(y_j,t) = -\frac{\alpha}{8}(4\bar{u}^2 - s^2) + f - \gamma. \tag{D12}$$

REFERENCES

[1] Barabasi A-L and H.E. Stanley, 1995, Fractal concepts in surface growth, Cambridge University, New York.

[2] Halpin-Healy,and Y.C.Zhang, Phys.Rep. **254**(1995); J. Krug, Adv. Phys. **46**, 139 (1997).

[3] J. Krug and H. Spohn (1990),In solids far from equilibrium growth,Morphology and defects, edited by C.Godreche(Cambridge University ,New York).

[4] P. Meakin Fractals,Scaling and Growth Far from Equilibrium, Cambridge Univ. Press, Cambridge (1998).

[5] Mehran Kardar, Physica A **281** 295 (2000).

[6] Edwards, S.F. and D.R. Wilkinson, 1982, Proc. R. Soc. London, Ser . A **381**,17.

[7] M. Kardar, G. Parisi and Y.C. Zhang, Phys. Rev. Lett. **56**, 889 (1986)

[8] F. Family and T. Vicsek, J. Phys. A **18**, L75 (1985)

[9] M. Marsilli, A. Maritan, F. Toigoend J.R. Banavar, Rev. of Modern Physics, **68**, 963 (1996)

[10] P. Meakin, P. Ramanlal,L.M. Sander and R.C. Ball, Phys. Rev. **A 34**, 5091 (1986)

[11] J. Krug, Phys. Rev. **A 36**, 5465 (1987)

[12] U. Dekker and F. Haake, Phys. Rev. **A11**, 2043 (1975).

[13] E. Frey and V.C. Tauber, Phys. Rev. **E50**, 1024 (1994)

[14] K.J. Wiese, Phys. Rev. **E56**, 5013 (1997); cond-mat/9802068.

[15] F. Colaiori and M.A. Moore, cond-mat/0010410

[16] E. Katzav and M. Schwartz, cond-mat/0012103

[17] L.H. Tang and H. Leschorn, Phys. Rev. A**45** 7162 (1992)

[18] T. Ala-Nissila, T. Hielt, J.M. Kosterlitz and O. Venalainen, J. Stat. Phys. **72** 207 (1993)

[19] J.M. Kim, Phys. Rev. lett **80**, 888 (1998)

[20] C.S. Chin and M. den Nijs, cond-mat/9810083

[21] M. Prahofer and H. Spohn, cond-mat/9912264

[22] M.K. Verma, Physica A**277** (2000) 359

[23] E. Marinari, A. Pagnani and G. Parisi, J. Phys. A: Math. Gen. 33 (2000) 8181, Cond-mat /0005105

[24] M.A. Moore et al, Phys.Rev.Lett, **74**, 4257 (1995)

[25] T.J. Newman and H. Kallabis, cond-mat/9512104

[26] T.J. Newman and A.J. Bray, cond-mat/9604071

[27] S. Stepanow, Phys. Rev. E **55**, R4853 (1997)

[28] M. Lassig, Nucl. Phys. B**448** (1998), cond-mat/9806330

[29] C. Castellano, M. Marsili and L. Pietronero, Phys. Rev. Lett. **80** (1998) 3525, cond-mat/9802284

[30] C. Castellano, A.Gabrielli, M. Marsili, M.A Munoz and L. Pietronero, Phys. Rev. E**58** (1998) R5209, cond-mat/9809197

[31] C. Castellano, M. Marsili, M.A. Munoz and L. Pietronero, cond-mat/9904434

[32] H.K. Janssen, U.C. Taeuber and E. Frey, Eur. Phys. J. **B 9** (1999) 491, cond-mat/9808325

[33] L. Golubovic and Z.G. Wang, Phys.Rev.**E49**, 2567 (1994).

[34] I. Ispolatov, P.L. Krapirsky and S. Redner Phys.Rev.**E52** ,2540 (1995).

[35] J. Krug and H. Spohn, Europhys Lett.**8**, 219 (1989).

[36] H. Van Beijeren, R. Kutner and H. Spohn, Phys.Rev.Lett**54**, 2026 (1985)

[37] H.k. Janssen and B. Schmittmann, Z. Phys.**B63**, 517 (1986).

[38] G. Blatter, M.V. Feigelman, V.B. Geshkenbein, A.I. Larkin and V.M. Vinokur, Rev. Modern Phys. **66** (1994) 1125

[39] D.A. Huse,C.l. Henley and D.S. Fisher,Phys.Rev.Lett.**55**, 2924(1985);

[40] M. Kardar and Y.C.Zhang,Phys.Rev.Lett.**58**,2087(1987)

[41] D.S. Fisher and D.A. Huse, Phys.Rev.**B43**,10728(1991).

[42] R.D. Kamien, P.Le. Doussal and D.R. Nelson, Phys. rev. A**45** (1992) 8727

[43] H.C. Fogedby, A.B. Eriksson and L.V. Mikheev, Phys. Rev. Lett. **75** (1995) 1883

[44] L. Balents, J-P. Bouchaud and M. Mezard, cond-mat/9601137

[45] R. Bundschuh and M. Lassig, cond-mat/9602045

[46] G. Parisi and F. Slanina, cond-mat/9712208

[47] D.A. Gorokhov and G. Blatter, Phys. Rev. lett. **82** (1999) 2705

[48] P. Delos Rios, Phys. Rev. Lett. **82** (1999) 4236

[49] A. Basu, Phys. Rev. E**62**, (2000) 4675

[50] E. Brunet and B. Derrida, Phys. rev. E **61**, (2000) 6789

[51] E. Perlsman and S. Havlin, Phys. Rev. E**63** (2001), cond-mat/0011157

[52] R. Mohayaee, A.L. stella and C. Vander Zande, cond-mat/0101091

[53] M.V. Feigelman, Sov. Phys. JETP **52**(1980) 555; translated from Zh. Eksper. Teoret. Fiz. **79**(1980),1095.

[54] S.F. Shandarin,Ya.B. Zeldovich, Rev. Modern Phys. **61** (1989) no, 2, 185

[55] M. Vergassola, B. Dubrulle, U. Frisch and A. Noullez, Astron. Astrophys, **280**, 325 (1994)

[56] S.F. Shandarin, astro-phy/9507082.

[57] U. Frisch, J. Bec and B. Villone, cond-mat/9912110.

[58] J. P. Bouchaud, M. Mezard, Phys. Rev. E **54**, 5116 (1996), cond-mat/9607006

[59] M. Mezard, cond-mat/9801029.

[60] H. C. Fogedby, Phys. Rev. E **57**, 2331 (1998); Phys. Rev. Let. **80**, 1126 (1998); Phys. Rev. E **57**, 4993 (1998); Phys. Rev. E **59**, 5065 (1999); Phys. Rev. E **66**, 4950 (1999).

[61] V. Yakhot and A. Chekhlov, Phys. Rev. Lett. **77** 3118 (1996)

[62] A.Chekhlov and V. Yakhot, Phys.Rev.E, **R**2739 (1995).

[63] A. Polyakov, Phys. Rev. E **52**, 6183 (1995).

[64] E. Balkovsky, G. Falkovich, I. Kolokolov and V. Lebedev, JETP Lett. **61**, 1012 (1995); Phys. Rev. Lett. **78** 1452 (1997); chao-dyn/9609005;

[65] G. Falkovich and V. Lebedev, chao-dyn/9708002.

[66] V. Gurarie and A. Migdal, Phys. Rev. E **54** 4908 (1996).

[67] J. P. Bouchaud, M. Mezard, Phys. Rev. E **54**, 5116 (1996), cond-mat/9607006.

[68] T. Gotoh and R. H. Kraichnan, chao-dyn/9803037.

[69] S. Boldyrev, Phys. Rev. E **55**, 6907 (1997), hep-th/9610080.

[70] S. Boldyrev, hep-th/9707255 ; hep-th/9805100.

[71] V. Yakhot, Phys. Rev. E **57**, 1737 (1997); chao-dyn/9904016; chao-dyn/9909017; chao-dyn/0001027.

[72] W.E, K.Khanin, A.Mazel, and Ya.G.Sinai, Phys.Rev.Lett., **78** , 1904 (1997).

[73] W. E and E. Vanden Eijnden, Phys.Rev.Lett.**83**,2572(1999) ;chao-dyn/9901006; chao-dyn/9904028; chao-dyn/9901029;

[74] R.H. Kraichnan, Phys. Fluids **11** (1999), 3738; chao-dyn/9901023.

[75] T. Gotoh and R.H. Kraichnan, Phys. Fluids A**5** (1993), 445 ; Phys. Fluids **10** (1998), 2859;

[76] J. Bec and U. Frisch, cond-mat/9906047; J. Bec, nlin.CD/0103029.

[77] J. Bec, U. Frisch, K. Khanin, chao-dyn/9910001.

[78] D. Bernard and K. Gawedzki, chao-dyn/9805002;

[79] Michel Bauer and Denis Bernard, chao-dyn/9812018.

[80] L. Frachebourg and Ph. A. Martin, cond-mat/9905056.

[81] S. N. Gurbatov, chao-dyn/9912011

[82] F. Hayot and C. Jayaprakash, nlin.CD/0005050;chao-dyn/9901026

[83] U. Frisch and J. Bec, Burgulence, to appear in Proceeding Les Houches 2000, New Trends in Turbulence, nlin-cd/0012033

[84] D. Bernard, cond-mat/0007106

[85] J. Davoudi, A.A. Masoudi, M.R. Rahimi Tabar, A.R. Rastegar and F. Shahbazi,Phys. Rev. E **63**, 056308 (2001)

[86] B. Derrida and J.L. Lebowitz, Phys.Rev.Lett.**80**,209(1998)

[87] B. Derrida and C. Appert, Statistical Physics,**94**,1,(1999)

[88] J.M. Kim, M.A. Moore and A.J. Bray, Phy. Rev. **A 44** (1991) 2345

[89] A.A. Masoudi, F. Shahbazi, J. Davoudi and M.R. Rahimi Tabar, in prepreation.

[90] J. Bec, nlin.CD/0103029

[91] M. Plischke etal. Phys. Rev. A **35**, 3485 (1987)

[92] J. Neergard and M. Den Nijs, J. Phys. A **30** 1935 (1997)

[93] J. Bec, Private communication.

Figure Captions

Figure.1 Behaviour of the time scales in which the moments $\langle (h-\tilde{h})^{2n} \rangle$ become negative in terms of n. The square points are calculated according to eq.(40) while the solid line is the fitting curve asymptotically tending to $\frac{1}{4}(\frac{k(0)}{\alpha^2 k_{xx}^2(0)})^{1/3}$.

Figure.2 In the upper graph the sharp valley solutions in KPZ equation are demonstrated while in lower one the corresponding shock structures in Burgers equation are sketched . The variables characterising the cusp, namely h_{x-} , h_{x+} and $\tilde{h}$ are shown.

Figure.3 Different time snapshots of gradient configuration whithin system size, i.e. $-\partial_x h$ verses x. The time scale for shock creation is demonstrated for $\sigma \sim L$. The solid points are showing the jumps in the height gradient. *J. Bec* [93] .

Figure.4 Different time snapshots of gradient configuration whithin system size, i.e. $-\partial_x h$ verses x. The time scale for shock creation is demonstrated for $\sigma < L$. The solid points are showing the jumps in the height gradient. *J. Bec* [93] .

Figure.5 Different time snapshots of gradient configuration whithin system size, i.e. $-\partial_x h$ verses x. The time scale for shock creation is demonstrated for $\sigma \ll L$. The solid points are showing the jumps in the height gradient. *J. Bec* [93].


t / t*
n
0
0.25
0.5
0.75
1
1.25
1.5
1.75
2
0
25
50
75
100


h(x)
x
h_{x-}
h_{x+}
$\tilde{h}$
$\bar{h}$
$-h_x(x)$
x



Fig.3



Fig.4



t=0.10

t=0.20

t=0.30

t=0.40

t=0.50

t=0.60

Fig.5